


RecomAb™

ERBB2 recombinant monoclonal antibody, clone SAIC-02A-7

Catalog # RAB03433 Size 200 ug


Applications



Immunofluorescence

Immunofluorescent staining of MCF-7 cells with ERBB2 recombinant monoclonal antibody, clone SAIC-02A-7 (Cat # RAB03433).

Immunofluorescence analysis of paraformaldehyde fixed MCF-7 cells stained with the chimeric r version of RAB03433 at 10 ug/mL followed by Alexa Fluor® 488 secondary antibody (2 ug/mL), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). The isotype control was stained with anti-Fluorescein antibody followed by staining with Alexa Fluor® 488 secondary antibody.



Flow Cytometry

Flow cytometric analysis of MCF-7 cells with ERBB2 recombinant monoclonal antibody, clone SAIC-02A-7 (Cat # RAB03433).

MCF-7 cells were stained with anti-Fluorescein IgG antibody (isotype control- black line) or the r1 version of RAB03433 (blue line) at a dilution of 1:100 for 1h at RT. After washing the bound antibody was detected using a goat anti-mouse IgG AlexaFluor® 488 antibody at a dilution of 1:1000 and cells analyzed using a FACSCanto flow-cytometer.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human ERBB2.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a peptide "GLQSLPTHDPSPQLQR" derived from the receptor tyrosine kinase erbB2 conjugated to KLH.
Reactivity	Human

Form	Liquid
Isotype	IgG
Recommend Usage	ELISA Flow Cytometry Immunofluorescence Immuno-MRM (multiple reaction monitoring) Western Blot The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS with 0.02% Proclin 300
Storage Instruction	Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

- Immunofluorescence

Immunofluorescent staining of MCF-7 cells with ERBB2 recombinant monoclonal antibody, clone SAIC-02A-7 (Cat # RAB03433).

Immunofluorescence analysis of paraformaldehyde fixed MCF-7 cells stained with the chimeric r version of RAB03433 at 10 ug/mL followed by Alexa Fluor® 488 secondary antibody (2 ug/mL), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). The isotype control was stained with anti-Fluorescein antibody followed by staining with Alexa Fluor® 488 secondary antibody.

- Enzyme-linked Immunoabsorbent Assay

- Flow Cytometry

Flow cytometric analysis of MCF-7 cells with ERBB2 recombinant monoclonal antibody, clone SAIC-02A-7 (Cat # RAB03433). MCF-7 cells were stained with anti-Fluorescein IgG antibody (isotype control-black line) or the r1 version of RAB03433 (blue line) at a dilution of 1:100 for 1h at RT. After washing the bound antibody was detected using a goat anti-mouse IgG AlexaFluor® 488 antibody at a dilution of 1:1000 and cells analyzed using a FACSCanto flow-cytometer.

- Immuno-MRM (multiple reaction monitoring)

Gene Info — ERBB2

Entrez GeneID [2064](#)

Gene Name ERBB2

Gene Alias	CD340, HER-2, HER-2/neu, HER2, NEU, NGL, TKR1
Gene Description	v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian)
Omim ID	137215 137800 164870 211980
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (position 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized. [provided by RefSeq]
Other Designations	c-erb B2/neu protein erbB-2 herstatin neuroblastoma/glioblastoma derived oncogene homolog v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived oncogene homolog)

Pathway

- [Adherens junction](#)
- [Bladder cancer](#)
- [Calcium signaling pathway](#)
- [Endometrial cancer](#)
- [ErbB signaling pathway](#)
- [Focal adhesion](#)
- [Non-small cell lung cancer](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)

Disease

- [Adenocarcinoma](#)
- [Ataxia telangiectasia](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cell Transformation](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Colorectal Neoplasms](#)
- [Disease Progression](#)
- [Endometrial Neoplasms](#)
- [Esophageal Neoplasms](#)
- [Fibroadenoma](#)
- [Gastritis](#)
- [Genetic Predisposition to Disease](#)
- [Glioma](#)
- [Head and Neck Neoplasms](#)
- [Heart Diseases](#)
- [Kidney Failure](#)
- [Laryngeal Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Mouth Neoplasms](#)

- [Neoplasm Invasiveness](#)
- [Neoplasm Metastasis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Obesity](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Papillomavirus Infections](#)
- [Pharyngeal Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Hyperplasia](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Skin Neoplasms](#)
- [Stomach Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Tooth Abnormalities](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Werner syndrome](#)