


RecomAb™

IL18 recombinant monoclonal antibody, clone SAIC-24B-4

Catalog # RAB03572 Size 200 ug


Applications



Western Blot

Western blot analysis of Lane 1: human peripheral blood monocyte (untreated) and Lane 2: human peripheral blood monocyte (treated with PMA/Ionomycin) with IL18 recombinant monoclonal antibody, clone SAIC-24B-4 (Cat # RAB03572).

Western Blot analysis the lysates (35ug protein in RIPA buffer) were resolved on a 15% SDS PAGE gel and blots were probed with the chimeric r version of RAB03572 at 0.3 ug/mL before detection using an anti-rondary antibody. Protein was detected by chemiluminescence.



Flow Cytometry

Flow cytometric analysis of human peripheral blood monocytes with IL18 recombinant monoclonal antibody, clone SAIC-24B-4 (Cat # RAB03572). The human peripheral blood monocytes were fixed using 2% PFA and permeabilized 0.5% Triton and stained with unimmunized r antibody antibody (isotype control-black line) or the chimeric r version of RAB03572 (blue line) at a dilution of 1:100 for 1h at RT. An additional population of PBMC's stimulated with PMA and Ionomycin in the presence of Monensin for 4 hours were stained in a similar manner (purple line). After washing the bound antibody was detected using a goat anti-mouse IgG AlexaFluor® 488 antibody at a dilution of 1:1000 and cells analyzed using a FACSCanto flow-cytometer.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human IL18.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against peptide "ISTLSCENK" derived from IL-18.

Reactivity	Human
Form	Liquid
Isotype	IgG
Recommend Usage	Flow Cytometry Immuno-MRM (multiple reaction monitoring) Western Blot The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.02% Proclin 300)
Storage Instruction	Store at 4°C for 3 months. For long term storage, aliquot and store at -20°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Western blot analysis of Lane 1: human peripheral blood monocyte (untreated) and Lane 2: human peripheral blood monocyte (treated with PMA/Ionomycin) with IL18 recombinant monoclonal antibody, clone SAIC-24B-4 (Cat # RAB03572).

Western Blot analysis the lysates (35ug protein in RIPA buffer) were resolved on a 15% SDS PAGE gel and blots were probed with the chimeric r version of RAB03572 at 0.3 ug/mL before detection using an anti-rondary antibody. Protein was detected by chemiluminescence.

- Flow Cytometry

Flow cytometric analysis of human peripheral blood monocytes with IL18 recombinant monoclonal antibody, clone SAIC-24B-4 (Cat # RAB03572).

The human peripheral blood monocytes were fixed using 2% PFA and permeabilized 0.5% Triton and stained with unimmunized r antibody antibody (isotype control-black line) or the chimeric r version of RAB03572 (blue line) at a dilution of 1:100 for 1h at RT. An additional population of PBMC's stimulated with PMA and Ionomycin in the presence of Monensin for 4 hours were stained in a similar manner (purple line). After washing the bound antibody was detected using a goat anti-mouse IgG AlexaFluor® 488 antibody at a dilution of 1:1000 and cells analyzed using a FACSCanto flow-cytometer.

- Immuno-MRM (multiple reaction monitoring)

Gene Info — IL18

Entrez GeneID	3606
Gene Name	IL18
Gene Alias	IGIF, IL-18, IL-1g, IL1F4, MGC12320
Gene Description	interleukin 18 (interferon-gamma-inducing factor)

Omim ID	600953
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a proinflammatory cytokine. This cytokine can induce the IFN-gamma production of T cells. The combination of this cytokine and IL12 has been shown to inhibit IL4 dependent IgE and IgG1 production, and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity. [provided by RefSeq]
Other Designations	IL-1 gamma interferon-gamma-inducing factor interleukin 18 interleukin-1 gamma interleukin-18

Pathway

- [Cytokine-cytokine receptor interaction](#)

Disease

- [Abortion](#)
- [Acute Lung Injury](#)
- [Adenocarcinoma](#)
- [Aggressive Periodontitis](#)
- [Alzheimer disease](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Atrial Fibrillation](#)
- [Autoimmune Diseases](#)
- [Behcet Syndrome](#)
- [Brain Ischemia](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)

- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Chorioamnionitis](#)
- [Choriocarcinoma](#)
- [Choroidal Neovascularization](#)
- [Chronic Periodontitis](#)
- [Cognition Disorders](#)
- [Colitis](#)
- [Colorectal Neoplasms](#)
- [Connective Tissue Diseases](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Crohn Disease](#)
- [Death](#)
- [Dermatitis](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Duodenal Ulcer](#)
- [Dyspepsia](#)
- [Edema](#)
- [Endometriosis](#)
- [Endotoxemia](#)
- [Epstein-Barr Virus Infections](#)

- [Esophageal Neoplasms](#)
- [Fetal Diseases](#)
- [Fetal Membranes](#)
- [Gastrointestinal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Giant Cell Arteritis](#)
- [Gingivitis](#)
- [Glomerulonephritis](#)
- [Graft vs Host Disease](#)
- [Graves Disease](#)
- [Graves Ophthalmopathy](#)
- [Head and Neck Neoplasms](#)
- [Heart Diseases](#)
- [Helicobacter Infections](#)
- [Hematologic Diseases](#)
- [Hemophilia A](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [HIV-Associated Lipodystrophy Syndrome](#)
- [Hodgkin Disease](#)
- [Hydatidiform Mole](#)
- [Hypersensitivity](#)
- [Hypertension](#)
- [Infant](#)
- [Infection](#)

- [Inflammation](#)
- [Inflammatory Bowel Diseases](#)
- [Insulin Resistance](#)
- [Intracranial Arteriosclerosis](#)
- [Kidney Calculi](#)
- [Kidney Failure](#)
- [Kidney Neoplasms](#)
- [Latex Hypersensitivity](#)
- [Leiomyoma](#)
- [Lichen Planus](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Lupus Nephritis](#)
- [Lymphadenitis](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Macular Degeneration](#)
- [Metabolic Syndrome X](#)
- [Metaplasia](#)
- [Mouth Neoplasms](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Musculoskeletal Diseases](#)
- [Mycobacterium Infections](#)
- [Myocardial Infarction](#)

- [Nasopharyngeal Neoplasms](#)
- [Neoplasms](#)
- [Obesity](#)
- [Obstetric Labor](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pain](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Periodontitis](#)
- [Pneumoconiosis](#)
- [Postoperative Complications](#)
- [Precancerous Conditions](#)
- [Pre-Eclampsia](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Proteinuria](#)
- [Pulmonary Disease](#)
- [Purpura](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Retinal Vein Occlusion](#)

- [Rhinitis](#)
- [Sarcoidosis](#)
- [Schistosomiasis mansoni](#)
- [Schizophrenia](#)
- [Skin Diseases](#)
- [Stomach Neoplasms](#)
- [Stomach Ulcer](#)
- [Stroke](#)
- [Substance Abuse](#)
- [Thyroid Diseases](#)
- [Tuberculosis](#)
- [Uterine Cervical Neoplasms](#)
- [Uterine Neoplasms](#)
- [Uveitis](#)
- [Vasculitis](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)
- [Wounds and Injuries](#)