


RecomAb™

CXCR2 recombinant monoclonal antibody, clone HC2

Catalog # RAB03596 Size 200 ug


Applications



Immunofluorescence

Immunofluorescent staining of human peripheral blood leukocytes (PBL) with CXCR2 recombinant monoclonal antibody, clone HC2 (Cat # RAB03596). Immunofluorescence analysis of paraformaldehyde fixed human PBL cells on Shi-fix™ coverslips stained with the chimeric r version of RAB03596 at 10 ug/mL for 1h followed by Alexa Fluor® 488 secondary antibody (2 ug/mL), showing membrane staining. The nuclear stain is DAPI (blue). The isotype control was an unknown specificity antibody followed by staining with Alexa Fluor® 488 secondary antibody.

- (A) RAB03596
- (B) DAPI
- (C) Merged channels
- (D) Isotype control



Flow Cytometry

Flow cytometric analysis of human peripheral blood monocytes (PBM) with CXCR2 recombinant monoclonal antibody, clone HC2 (Cat # RAB03596). Paraformaldehyde fixed human PBM cells were stained with anti-unknown specificity antibody (isotype control-black line) or the r version of RAB03596 (blue line) at a dilution of 1:100 for 1h at RT. After washing the bound antibody was detected using a goat anti-r AlexaFluor® 488 antibody at a dilution of 1:1000 and cells analyzed using a FACSCanto flow-cytometer.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human CXCR2.
Antibody Species	Rabbit

Immunogen	Original antibody is raised against human CXCR2.
Reactivity	Human
Form	Liquid
Isotype	IgG
Recommend Usage	Flow Cytometry Immunofluorescence The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.02% Proclin 300)
Storage Instruction	Store at 4°C for 3 months. For long term storage, aliquot and store at -20°C. Aliquot to avoid repeated freezing and thawing.

Applications

● Immunofluorescence

Immunofluorescent staining of human peripheral blood leukocytes (PBL) with CXCR2 recombinant monoclonal antibody, clone HC2 (Cat # RAB03596).

Immunofluorescence analysis of paraformaldehyde fixed human PBL cells on Shi-fix™ coverslips stained with the chimeric r version of RAB03596 at 10 ug/mL for 1h followed by Alexa Fluor® 488 secondary antibody (2 ug/mL), showing membrane staining. The nuclear stain is DAPI (blue). The isotype control was an unknown specificity antibody followed by staining with Alexa Fluor® 488 secondary antibody.

(A) RAB03596

(B) DAPI

(C) Merged channels

(D) Isotype control

● Flow Cytometry

Flow cytometric analysis of human peripheral blood monocytes (PBM) with CXCR2 recombinant monoclonal antibody, clone HC2 (Cat # RAB03596).

Paraformaldehyde fixed human PBM cells were stained with anti-unknown specificity antibody (isotype control-black line) or the r version of RAB03596 (blue line) at a dilution of 1:100 for 1h at RT. After washing the bound antibody was detected using a goat anti-r AlexaFluor® 488 antibody at a dilution of 1:1000 and cells analyzed using a FACSCanto flow-cytometer.

Gene Info — CXCR2

Entrez GeneID	3579
Gene Name	CXCR2
Gene Alias	CD182, CDw128b, CMKAR2, IL8R2, IL8RA, IL8RB

Gene Description	chemokine (C-X-C motif) receptor 2
Omim ID	146928
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. This gene, IL8RA, a gene encoding another high affinity IL8 receptor, as well as IL8RBP, a pseudogene of IL8RB, form a gene cluster in a region mapped to chromosome 2q33-q36. [provided by RefSeq]
Other Designations	C-X-C chemokine receptor type 2 CXC-R2 CXCR-2 CXCR2 gene for IL8 receptor type B GRO/MGSA receptor IL-8 receptor type 2 IL-8RB chemokine (CXC) receptor 2 high affinity interleukin-8 receptor B interleukin 8 receptor B interleukin 8 receptor type 2 interl

Pathway

- [Chemokine signaling pathway](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)

Disease

- [Acquired Immunodeficiency Syndrome](#)
- [Adenocarcinoma](#)
- [Alcoholism](#)
- [Asthma](#)
- [Bacteriuria](#)
- [Behcet Syndrome](#)
- [Biliary Tract Neoplasms](#)

- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Conduct Disorder](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Helicobacter Infections](#)
- [Hematologic Diseases](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Infant](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Leishmaniasis](#)

- [Leukemia](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Maduromycosis](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Multiple Sclerosis](#)
- [Neoplasm Invasiveness](#)
- [Obstetric Labor](#)
- [Occupational Diseases](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Papillomavirus Infections](#)
- [Periodontal Diseases](#)
- [Periodontitis](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoma](#)
- [Stomach Neoplasms](#)

- [Tuberculosis](#)
- [Uterine Cervical Neoplasms](#)
- [Viremia](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)