


RecomAb™

CDK4 recombinant monoclonal antibody, clone 8H4

Catalog # RAB04018 Size 100 uL

Applications



Western Blot

Western Blot analysis of Lane 1: Jurkat whole cell lysate; Lane 2: HeLa whole cell lysate; Lane 3: 293 whole cell lysate; Lane 4: MCF-7 whole cell lysate; Lane 5: HepG2 whole cell lysate; Lane 6: K562 whole cell lysate; Lane 7: A549 whole cell lysate.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human CDK4.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against recombinant protein corresponding to full length human CDK4.
Reactivity	Human
Form	Liquid
Purification	Affinity-chromatography
Isotype	IgG
Recommend Usage	ELISA Western Blot (1:500-1:5000) The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS, pH7.4 (150mM NaCl, 50% glycerol and 0.02% sodium azide)
Storage Instruction	store at -20 °C or -80 °C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot

Western Blot analysis of Lane 1: Jurkat whole cell lysate; Lane 2: Hela whole cell lysate; Lane 3: 293 whole cell lysate; Lane 4: MCF-7 whole cell lysate; Lane 5: HepG2 whole cell lysate; Lane 6: K562 whole cell lysate; Lane 7: A549 whole cell lysate.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — CDK4

Entrez GeneID	1019
Protein Accession#	P11802
Gene Name	CDK4
Gene Alias	CMM3, MGC14458, PSK-J3
Gene Description	cyclin-dependent kinase 4
Omim ID	123829 609048
Gene Ontology	Hyperlink

Gene Summary	The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is highly similar to the gene products of <i>S. cerevisiae</i> cdc28 and <i>S. pombe</i> cdc2. It is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression. The activity of this kinase is restricted to the G1-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p16(INK4a). This kinase was shown to be responsible for the phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and Rb were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation sites of this gene have been reported. [provided by RefSeq]
--------------	--

Other Designations	cell division kinase 4 melanoma cutaneous malignant, 3
--------------------	--

Pathway

- [Bladder cancer](#)
- [Cell cycle](#)
- [Chronic myeloid leukemia](#)
- [Glioma](#)
- [Melanoma](#)
- [Non-small cell lung cancer](#)
- [p53 signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)
- [T cell receptor signaling pathway](#)
- [Tight junction](#)

Disease

- [Adenocarcinoma](#)
- [Birth Weight](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)

- [Glioblastoma](#)
- [Head and Neck Neoplasms](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Mouth Neoplasms](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Obesity](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Precancerous Conditions](#)
- [Pulmonary Disease](#)
- [Retinoblastoma](#)
- [Skin Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Uveal Neoplasms](#)
- [Werner syndrome](#)