


RecomAb™

CXCR4 recombinant monoclonal antibody, clone 2F10

Catalog # RAB04048 Size 100 uL

Applications



Immunofluorescence

Immunofluorescence staining of HeLa Cells with CXCR4 recombinant monoclonal antibody, clone 2F10 at 1:50, counter-stained with DAPI.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human CXCR4.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against recombinant protein corresponding to full length human CXCR4.
Reactivity	Human
Form	Liquid
Purification	Affinity-chromatography
Isotype	IgG
Recommend Usage	ELISA Flow Cytometry (1:20-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH7.4 (150mM NaCl, 50% glycerol and 0.02% sodium azide)
Storage Instruction	Store at -20 °C or -80 °C. Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunofluorescence

Immunofluorescence staining of Hela Cells with CXCR4 recombinant monoclonal antibody, clone 2F10 at 1:50, counter-stained with DAPI.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — CXCR4

Entrez GeneID	7852
Protein Accession#	P61073
Gene Name	CXCR4
Gene Alias	CD184, D2S201E, FB22, HM89, HSY3RR, LAP3, LCR1, LESTR, NPY3R, NPYR, NPYRL, NPY3R, WHIM
Gene Description	chemokine (C-X-C motif) receptor 4
Omim ID	162643 193670
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a CXC chemokine receptor specific for stromal cell-derived factor-1. The protein has 7 transmembrane regions and is located on the cell surface. It acts with the CD4 protein to support HIV entry into cells and is also highly expressed in breast cancer cells. Mutations in this gene have been associated with WHIM (warts, hypogammaglobulinemia, infections, and myelodysplasia) syndrome. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq]
Other Designations	C-X-C chemokine receptor type 4 CD184 antigen chemokine (C-X-C motif), receptor 4 (fusin) chemokine receptor 4 fusin leukocyte-derived seven-transmembrane-domain receptor lipopolysaccharide-associated protein 3 neuropeptide Y receptor Y3 seven transmembrane

Pathway

- [Axon guidance](#)

- [Chemokine signaling pathway](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Leukocyte transendothelial migration](#)

Disease

- [Acquired Immunodeficiency Syndrome](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Colon cancer](#)
- [Colorectal Neoplasms](#)
- [Coronary Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Growth Hormone-Secreting Pituitary Adenoma](#)
- [Hepatitis B](#)
- [HIV Infections](#)
- [Hyperglycemia](#)
- [Liver Neoplasms](#)

- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Mouth Neoplasms](#)
- [Multiple Sclerosis](#)
- [Neoplasm Invasiveness](#)
- [Pituitary Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Tumor Virus Infections](#)