


RecomAb™

HSPA5 recombinant monoclonal antibody, clone 4H12

Catalog # RAB04146 Size 100 uL

Applications



Immunohistochemistry

Immunohistochemistry image of HSPA5 recombinant monoclonal antibody, clone 4H12 diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human HSPA5.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against recombinant protein corresponding to full length human HSPA5.
Reactivity	Human
Form	Liquid
Purification	Affinity-chromatography
Isotype	IgG
Recommend Usage	ELISA Immunohistochemistry (1:50-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH7.4 (150 mM NaCl, 50% glycerol and 0.02% sodium azide)
Storage Instruction	Store at -20°C or -80°C. Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry

Immunohistochemistry image of HSPA5 recombinant monoclonal antibody, clone 4H12 diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — HSPA5

Entrez GeneID [3309](#)

Protein Accession# [P11021](#)

Gene Name HSPA5

Gene Alias BIP, FLJ26106, GRP78, MIF2

Gene Description heat shock 70kDa protein 5 (glucose-regulated protein, 78kDa)

Omim ID [138120](#)

Gene Ontology [Hyperlink](#)

Gene Summary When Chinese hamster K12 cells are starved of glucose, the synthesis of several proteins, called glucose-regulated proteins (GRPs), is markedly increased. Hendershot et al. (1994) [PubMed 80 20977] pointed out that one of these, GRP78 (HSPA5), also referred to as 'immunoglobulin heavy chain-binding protein' (BiP), is a member of the heat-shock protein-70 (HSP70) family and is involved in the folding and assembly of proteins in the endoplasmic reticulum (ER). Because so many ER proteins interact transiently with GRP78, it may play a key role in monitoring protein transport through the cell.[supplied by OMIM]

Other Designations Heat-shock 70kD protein-5 (glucose-regulated protein, 78kD)|OTTHUMP00000022124|heat shock 70kD protein 5 (glucose-regulated protein, 78kD)|heat shock 70kDa protein 5

Pathway

- [Antigen processing and presentation](#)

- [Prion diseases](#)

Disease

- [Alzheimer disease](#)
- [Bipolar Disorder](#)
- [Carcinoma](#)
- [Cognition](#)
- [Disease Susceptibility](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis B](#)
- [Infection](#)
- [Kidney Failure](#)
- [Liver Neoplasms](#)
- [Parkinson disease](#)
- [Schizophrenia](#)
- [Stress](#)