
#### RecomAb™

# MAPK3 (phospho T202) /MAPK1 (phospho T185) recombinant monoclonal antibody, clone 2D5

Catalog # RAB04304 Size 100 uL

# Applications




#### Western Blot

Western blot analysis of Lane 1:A549 whole cell lysate (treated with EGF 100ng/ml/20mins), Lane 2: A549 whole cell lysate (not treated), Lane 3: HepG2 whole cell lysate (treated with EGF 100ng/ml/20mins) and Lane 4: HepG2 whole cell lysate (not treated) with MAPK3 (phospho T202) +MAPK1 (phospho T185) recombinant monoclonal antibody, clone 2D5 (Cat # RAB04304).

#### Immunohistochemistry

Immunohistochemical staining of human glioma cancer with MAPK3 (phospho T202) +MAPK1 (phospho T185) recombinant monoclonal antibody, clone 2D5 (Cat # RAB04304) (diluated at 1:100)



#### Immunofluorescence

Immunofluorescent staining of Hela cells with MAPK3 (phospho T202) +MAPK1 (phospho T185) recombinant monoclonal antibody, clone 2D5 (Cat # RAB04304) (diluated at 1:100). The secondary antibody was Alexa Fluor 488congugated goat anti-rabbit IgG (green). Counter-stain DAPI was used (blue). 😵 Abnova

# **Product Information**

Product Description	Rabbit recombinant monoclonal antibody raised against human MAPK3.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic phosphopeptide corresponding to residues surroundin g T202 of human MAPK3 and T185 of human MAPK1.
Theoretical MW (kDa)	Calculated MW: 42 kD
Reactivity	Human
Form	Liquid
Purification	Affinity chromatography
lsotype	lgG
Recommend Usage	ELISA
	Immunofluorescence (1:20-1:200)
	Immunohistochemistry (1:50-1:200)
	Western Blot (1:500-1:5000)
	The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCI, pH 7.4 (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20 °C or -80 °C.
	Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul
	d be handled by trained staff only.

### **Applications**

Western Blot

Western blot analysis of Lane 1:A549 whole cell lysate (treated with EGF 100ng/ml/20mins), Lane 2: A549 whole cell lysate (not treated), Lane 3: HepG2 whole cell lysate (treated with EGF 100ng/ml/20mins) and Lane 4: HepG2 whole cell lysate (not treated) with MAPK3 (phospho T202) +MAPK1 (phospho T185) recombinant monoclonal antibody, clone 2D5 (Cat # RAB04304).

Immunohistochemistry

Immunohistochemical staining of human glioma cancer with MAPK3 (phospho T202) +MAPK1 (phospho T185) recombinant monoclonal antibody, clone 2D5 (Cat # RAB04304) (diluated at 1:100)

#### Immunofluorescence

Immunofluorescent staining of Hela cells with MAPK3 (phospho T202) +MAPK1 (phospho T185) recombinant monoclonal antibody, clone 2D5 (Cat # RAB04304) (diluated at 1:100). The secondary antibody was Alexa Fluor 488-congugated goat anti-rabbit IgG (green). Counter-stain DAPI was used (blue).

• Enzyme-linked Immunoabsorbent Assay

Gene Info — MAPK3	
Entrez GenelD	<u>5595</u>
Protein Accession#	<u>P27361</u>
Gene Name	MAPK3
Gene Alias	ERK1, HS44KDAP, HUMKER1A, MGC20180, P44ERK1, P44MAPK, PRKM3
Gene Description	mitogen-activated protein kinase 3
Omim ID	<u>601795</u>
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also kno wn as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates vari ous cellular processes such as proliferation, differentiation, and cell cycle progression in respons e to a variety of extracellular signals. This kinase is activated by upstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcrip t variants encoding different protein isoforms have been described. [provided by RefSeq
Other Designations	OTTHUMP00000174538 OTTHUMP00000174540 extracellular signal-regulated kinase 1 extrace llular signal-related kinase 1

# Pathway

- Acute myeloid leukemia
- <u>Adherens junction</u>
- Axon guidance
- <u>B cell receptor signaling pathway</u>
- Bladder cancer
- Chemokine signaling pathway
- Chronic myeloid leukemia
- Colorectal cancer

# 😵 Abnova

- Dorso-ventral axis formation
- Endometrial cancer
- ErbB signaling pathway
- Fc epsilon RI signaling pathway
- Fc gamma R-mediated phagocytosis
- Focal adhesion
- Gap junction
- Glioma
- <u>GnRH signaling pathway</u>
- Insulin signaling pathway
- Long-term depression
- Long-term potentiation
- MAPK signaling pathway
- Melanogenesis
- <u>Melanoma</u>
- mTOR signaling pathway
- <u>Natural killer cell mediated cytotoxicity</u>
- <u>Neurotrophin signaling pathway</u>
- Non-small cell lung cancer
- Pancreatic cancer
- Pathways in cancer
- Prion diseases
- Prostate cancer
- <u>Regulation of actin cytoskeleton</u>
- Renal cell carcinoma
- <u>T cell receptor signaling pathway</u>

# 😵 Abnova

- TGF-beta signaling pathway
- Thyroid cancer
- Toll-like receptor signaling pathway
- Type II diabetes mellitus
- <u>Vascular smooth muscle contraction</u>
- VEGF signaling pathway

## Disease

- Asthma
- <u>Autistic Disorder</u>
- Disease Models
- Genetic Predisposition to Disease