


RecomAb™

UBE4B recombinant monoclonal antibody, clone R04-6C6

Catalog # RAB04421 Size 100 uL

Applications



Western Blot

Western blot analysis of Lane 1: PC-12 whole cell lysate and Lane 2: C6 whole cell lysate with UBE4B recombinant monoclonal antibody, clone R04-6C6 (Cat # RAB04421).

| Specification | |
|----------------------|---|
| Product Description | Rabbit recombinant monoclonal antibody raised against human UBE4B. |
| Antibody Species | Rabbit |
| Immunogen | Original antibody is raised against a synthetic peptide corresponding to human UBE4B. |
| Theoretical MW (kDa) | Calculated MW: 146 k |
| Reactivity | Human, Rat |
| Form | Liquid |
| Purification | Affinity purification |
| lsotype | lgG |
| Recommend Usage | Immunocytochemistry Immunofluorescence (1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user. |
| Storage Buffer | In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA) |

Copyright © 2023 Abnova Corporation. All Rights Reserved.

😵 Abnova

Product Information

Storage Instruction

Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot

Western blot analysis of Lane 1: PC-12 whole cell lysate and Lane 2: C6 whole cell lysate with UBE4B recombinant monoclonal antibody, clone R04-6C6 (Cat # RAB04421).

- Immunocytochemistry
- Immunofluorescence

Gene Info — UBE4B

| Entrez GenelD | <u>10277</u> |
|--------------------|---|
| Protein Accession# | <u>O95155</u> |
| Gene Name | UBE4B |
| Gene Alias | E4, HDNB1, KIAA0684, UBOX3, UFD2 |
| Gene Description | ubiquitination factor E4B (UFD2 homolog, yeast) |
| Gene Ontology | Hyperlink |
| Gene Summary | The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnor mal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzym es: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-prot ein ligases, or E3s. This gene encodes an additional conjugation factor, E4, which is involved in multiubiquitin chain assembly. This gene is also the strongest candidate in the neuroblastoma tum or suppressor genes. Alternatively spliced transcript variants encoding distinct isoforms have bee n found for this gene. [provided by RefSeq |
| Other Designations | OTTHUMP00000001726 OTTHUMP00000001727 homologous to yeast UFD2 homozygously del eted in neuroblastoma-1 ubiquitin-fusion degradation protein 2 ubiquitination factor E4B ubiquitin ation factor E4B (homologous to yeast UFD2) |

Pathway

Copyright © 2023 Abnova Corporation. All Rights Reserved.



<u>Ubiquitin mediated proteolysis</u>

Disease

- <u>Carcinoma</u>
- Genetic Predisposition to Disease
- Hepatitis B
- Liver Neoplasms