


RecomAb™

STAT6 recombinant monoclonal antibody, clone R04-0B5

Catalog # RAB04439 Size 100 uL

Applications



Western Blot

Western blot analysis of Lane 1: HeLa whole cell lysate and Lane 2: U2OS whole cell lysate with STAT6 recombinant monoclonal antibody, clone R04-0B5 (Cat # RAB04439).

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human STAT6.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human STAT6.
Theoretical MW (kDa)	Calculated MW: 94 kD
Reactivity	Human
Form	Liquid
Purification	Affinity chromatography
Isotype	IgG
Recommend Usage	Immunoprecipitation (1:20) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 50mM Tris-Glycine, 150mM NaCl, pH 7.4 (40% glycerol, 0.05% BSA and 0.01% Sodium azide)

Storage Instruction

Store at 4°C. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot

Western blot analysis of Lane 1: HeLa whole cell lysate and Lane 2: U2OS whole cell lysate with STAT6 recombinant monoclonal antibody, clone R04-0B5 (Cat # RAB04439).

- Immunoprecipitation

Gene Info — STAT6

Entrez GeneID[6778](#)**Protein Accession#**[P42226](#)**Gene Name**

STAT6

Gene Alias

D12S1644, IL-4-STAT, STAT6B, STAT6C

Gene Description

signal transducer and activator of transcription 6, interleukin-4 induced

Omim ID[601512](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4. Knockout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expression of cell surface markers, and class switch of immunoglobulins. [provided by RefSeq]

Other Designations

STAT, interleukin4-induced|signal transducer and activator of transcription 6|transcription factor IL-4 STAT

Pathway

- [Jak-STAT signaling pathway](#)

Disease

- [Anaphylaxis](#)
- [Arthritis](#)
- [Ascariasis](#)
- [Asthma](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Colitis](#)
- [Confusion](#)
- [Crohn Disease](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Drug Hypersensitivity](#)
- [Edema](#)
- [Endometriosis](#)
- [Food Hypersensitivity](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)

- [Graves Disease](#)
- [Hypersensitivity](#)
- [Infant](#)
- [Infertility](#)
- [Inflammatory Bowel Diseases](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Liver Diseases](#)
- [Lupus Erythematosus](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Nausea](#)
- [Neoplasms](#)
- [Nephrosis](#)
- [Nephrotic Syndrome](#)
- [Nut Hypersensitivity](#)
- [Pain](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)