


RecomAb™

P4HB recombinant monoclonal antibody, clone R08-6I8

Catalog # RAB04892 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of K562, C6 lysates with P4HB recombinant monoclonal antibody, clone R08-6I8 (Cat # RAB04892).

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human P4HB.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human P4HB.
Theoretical MW (kDa)	57
Reactivity	Human, Mouse, Rat
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 50 mM Tris-Glycine, pH 7.4 (0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction

Store at 4°C. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of K562, C6 lysates with P4HB recombinant monoclonal antibody, clone R08-6I8 (Cat # RAB04892).

Gene Info — P4HB

Entrez GeneID[5034](#)**Protein Accession#**[P07237](#)**Gene Name**

P4HB

Gene Alias

DSI, ERBA2L, GIT, P4Hbeta, PDI, PDIA1, PHDB, PO4DB, PO4HB, PROHB

Gene Description

prolyl 4-hydroxylase, beta polypeptide

Omim ID[176790](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes the beta subunit of prolyl 4-hydroxylase, a highly abundant multifunctional enzyme that belongs to the protein disulfide isomerase family. When present as a tetramer consisting of two alpha and two beta subunits, this enzyme is involved in hydroxylation of prolyl residues in procollagen. This enzyme is also a disulfide isomerase containing two thioredoxin domains that catalyze the formation, breakage and rearrangement of disulfide bonds. Other known functions include its ability to act as a chaperone that inhibits aggregation of misfolded proteins in a concentration-dependent manner, its ability to bind thyroid hormone, its role in both the influx and efflux of S-nitrosothiol-bound nitric oxide, and its function as a subunit of the microsomal triglyceride transfer protein complex. [provided by RefSeq]

Other Designations

collagen prolyl 4-hydroxylase beta|glutathione-insulin transhydrogenase|procollagen-proline, 2-oxo glutarate 4-dioxygenase (proline 4-hydroxylase), beta polypeptide|prolyl 4-hydroxylase, beta subunit|protein disulfide isomerase family A, member 1|protein d

Disease

- [Cardiovascular Diseases](#)

- [Diabetes Mellitus](#)
- [Edema](#)