


RecomAb™

PDGFRA recombinant monoclonal antibody, clone R06-6D3

Catalog # RAB04893 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of C6, 3T3 lysates with PDGFRA recombinant monoclonal antibody, clone R06-6D3 (Cat # RAB04893).

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human PDGFRA.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human PDGFRA.
Theoretical MW (kDa)	Calculated MW: 123 k
Reactivity	Human, Mouse, Rat
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 50 mM Tris-Glycine, pH 7.4 (0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction

Store at 4°C. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of C6, 3T3 lysates with PDGFRA recombinant monoclonal antibody, clone R06-6D3 (Cat # RAB04893).

Gene Info — PDGFRA

Entrez GeneID[5156](#)**Protein Accession#**[P16234](#)**Gene Name**

PDGFRA

Gene Alias

CD140A, MGC74795, PDGFR2, Rhe-PDGFRA

Gene Description

platelet-derived growth factor receptor, alpha polypeptide

Omim ID[173490](#) [606764](#) [607685](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. Studies in knockout mice, where homozygosity is lethal, indicate that the alpha form of the platelet-derived growth factor receptor is particularly important for kidney development since mice heterozygous for the receptor exhibit defective kidney phenotypes. [provided by RefSeq]

Other Designations

FIP1L1/PDGFRA fusion protein|platelet-derived growth factor receptor alpha|rearranged-in-hyper eosinophilia-platelet derived growth factor receptor alpha fusion protein

Pathway

- [Calcium signaling pathway](#)
- [Colorectal cancer](#)

- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Focal adhesion](#)
- [Gap junction](#)
- [Glioma](#)
- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Acute Disease](#)
- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Aneuploidy](#)
- [Asthma](#)
- [Brain Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)

- [Disease Progression](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Gastrointestinal Stromal Tumors](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Hyperparathyroidism](#)
- [Leukemia](#)
- [Liver Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Malignant melanoma](#)
- [Meningomyelocele](#)
- [Metabolic Syndrome X](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neural Tube Defects](#)
- [Osteoporosis](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Spinal Dysraphism](#)
- [Subdural Effusion](#)
- [Tobacco Use Disorder](#)
- [Tooth Abnormalities](#)
- [Uterine Cervical Neoplasms](#)
- [Vitiligo](#)