

RecomAb™

HNRNPA1 recombinant monoclonal antibody, clone R01-2A9

Catalog # RAB06475 Size 100 uL

Product Description Rabbit recombinant monoclonal antibody raised against human HNRNPA1.		
Antibody Species Rabbit Immunogen Original antibody is raised against recombinant protein corresponding to human HNRNPA1. Theoretical MW (kDa) Calculated MW: 39 kD Reactivity Human, Mouse, Rat Form Liquid Purification Affinity purification Isotype IgG Recommend Usage Flow Cytometry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Specification	
Immunogen Original antibody is raised against recombinant protein corresponding to human HNRNPA1. Theoretical MW (kDa) Calculated MW: 39 kD Reactivity Human, Mouse, Rat Form Liquid Purification Affinity purification Isotype IgG Recommend Usage Flow Cytometry (1:50-1:100) Immunofistochemistry (1:50-1:100) Immunofistochemistry (1:50-1:200) Western Blot (1:50-1:200) Western Blot (1:50-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Product Description	Rabbit recombinant monoclonal antibody raised against human HNRNPA1.
Theoretical MW (kDa) Calculated MW: 39 kD Reactivity Human, Mouse, Rat Form Liquid Purification Affinity purification Isotype IgG Recommend Usage Flow Cytometry (1:50-1:100) Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Antibody Species	Rabbit
Reactivity Human, Mouse, Rat Form Liquid Purification Affinity purification Isotype IgG Recommend Usage Flow Cytometry (1:50-1:100) Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Immunogen	Original antibody is raised against recombinant protein corresponding to human HNRNPA1.
Form Liquid Purification Affinity purification Isotype IgG Recommend Usage Flow Cytometry (1:50-1:100) Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Theoretical MW (kDa)	Calculated MW: 39 kD
Purification Affinity purification Isotype IgG Recommend Usage Flow Cytometry (1:50-1:100) Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Reactivity	Human, Mouse, Rat
Recommend Usage Flow Cytometry (1:50-1:100) Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Form	Liquid
Recommend Usage Flow Cytometry (1:50-1:100) Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Purification	Affinity purification
Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide) Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Isotype	lgG
Storage Instruction Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.	Recommend Usage	Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Western Blot (1:500-1:1000)
Aliquot to avoid repeated freezing and thawing.	Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide)
Note This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which sho	Storage Instruction	
d be handled by trained staff only.	Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot



- Immunohistochemistry
- Immunofluorescence
- Flow Cytometry

Gene Info — HNRNPA1	
Entrez GeneID	3178
Gene Name	HNRNPA1
Gene Alias	HNRPA1, MGC102835
Gene Description	heterogeneous nuclear ribonucleoprotein A1
Omim ID	<u>164017</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene belongs to the A/B subfamily of ubiquitously expressed heterogeneous nuclear ribonucl eoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind to RNAs. It is one of the most abundant core proteins of hnRNP complexes and it is localized to the nucleoplasm. This protein, along with other hnRNP proteins, is exported from the nucleus, probably bound to mRNA, and is immediately re-imported. Its M9 domain acts as both a nuclear localization and nuclear export signal. The encoded protein is involved in the packaging of pre-mRNA into hnRNP particles, transport of poly A+ mRNA from the nucleus to the cytoplasm, and may modulate splice site selection. It is als o thought have a primary role in the formation of specific myometrial protein species in parturition. Multiple alternatively spliced transcript variants have been found for this gene but only two transcripts are fully described. These variants have multiple alternative transcription initiation sites and multiple polyA sites. [provided by RefSeq
Other Designations	helix-destabilizing protein heterogeneous nuclear ribonucleoprotein A1B protein heterogeneous nuclear ribonucleoprotein B2 protein heterogeneous nuclear ribonucleoprotein core protein A1 nuclear ribonucleoprotein particle A1 protein single-strand DNA-bind

Disease

- Cardiovascular Diseases
- Diabetes Mellitus



• Edema