


RecomAb™

## MME recombinant monoclonal antibody, clone 10G11

Catalog # RAB07412 Size 100 uL

### **Applications**



# Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical analysis of paraffin-embedded human kidney tissue using MME recombinant monoclonal antibody, clone 10G11 (Cat # RAB07412) on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Specification	
Product Description	Rabbit recombinant monoclonal antibody raised against human MME.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human MME.
Reactivity	Human
Form	Liquid
Purification	Affinity chromatography purification
Isotype	lgG
Recommend Usage	ELISA Immunohistochemistry (1:50-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH7.4 (150 mM NaCl, 0.02% sodium azide and 50% glycerol)



#### **Product Information**

Storage Instruction	Store at -20°C or -80°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## **Applications**

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human kidney tissue using MME recombinant monoclonal antibody, clone 10G11 (Cat # RAB07412) on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Enzyme-linked Immunoabsorbent Assay

Gene Info — MME	
Entrez GenelD	4311
Protein Accession#	P08473
Gene Name	MME
Gene Alias	CALLA, CD10, DKFZp686O16152, MGC126681, MGC126707, NEP
Gene Description	membrane metallo-endopeptidase
Omim ID	120520
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a common acute lymphocytic leukemia antigen that is an important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on I eukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney, where it is present on the brush border of proximal tubules and on glomerular epithelium. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. This gene, which encodes a 100-k D type II transmembrane glycoprotein, exists in a single copy of greater than 45 kb. The 5' untranslated region of this gene is alternatively spliced, resulting in four separate mRNA transcripts. The coding region is not affected by alternative splicing. [provided by RefSeq



#### **Product Information**

**Other Designations** 

atriopeptidase|common acute lymphocytic leukemia antigen|enkephalinase|membrane metallo-en dopeptidase (neutral endopeptidase, enkephalinase)|membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10)|membrane metallo-endopeptidase

### Pathway

- Hematopoietic cell lineage
- Renin-angiotensin system

#### Disease

- Alzheimer disease
- Anorexia Nervosa
- Atherosclerosis
- Brain Injuries
- Bulimia
- Calcinosis
- Cardiovascular Diseases
- Coronary Artery Disease
- Diabetes Complications
- Genetic Predisposition to Disease
- Metabolic Syndrome X
- Neoplasms
- Osteoporosis