

## Datasheet

### CASP1 monoclonal antibody (M01), clone 3D2

**Catalog Number:** H00000834-M01

**Regulatory Status:** For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against a partial recombinant CASP1.

**Clone Name:** 3D2

**Immunogen:** CASP1 (AAH62327, 1 a.a. ~ 100 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Sequence:**

MADKVLKEKRKLFIRSMGEAPQAVQDNPAMPTSSGSE  
GNVKLCSLEEAQRIWKQKSAEIYPIMDKSSRTRLALIIC  
NEEFDSIPRRTGAEVDITGMTLL

**Host:** Mouse

**Reactivity:** Human

**Applications:** ELISA, IHC-P, S-ELISA, WB-Re, WB-Tr  
(See our web site product page for detailed applications information)

**Protocols:** See our web site at  
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Isotype:** IgG2a Kappa

**Storage Buffer:** In 1x PBS, pH 7.4

**Storage Instruction:** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 834

**Gene Symbol:** CASP1

**Gene Alias:** ICE, IL1BC, P45

**Gene Summary:** This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell

apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. This gene was identified by its ability to proteolytically cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes such as inflammation, septic shock, and wound healing. This gene has been shown to induce cell apoptosis and may function in various developmental stages. Studies of a similar gene in mouse suggest a role in the pathogenesis of Huntington disease. Alternative splicing of this gene results in five transcript variants encoding distinct isoforms. [provided by RefSeq]