
Bioactive

IL15 (Human) Recombinant Protein

Catalog # P3462 Size 50 ug

Applications



Specification

Product Description	Human IL15 (NP_751914, 49 a.a. - 162 a.a.) partial recombinant protein with His tag expressed in *E scherichia coli*.
Sequence	MNWVNVISDLKKIEDLIQSMHIDATLYTESDVHPSCKVTAMKCFLLELQVISLESGDASIHDTVENLII LANNSLSSNGNVTESGCKECEELEEKNIKEFLQSFVHIVQMFINTSLEHHHHHH
Host	Escherichia coli
Theoretical MW (kDa)	13.9
Form	Liquid
Preparation Method	*Escherichia coli* expression system
Purification	Conventional Chromatography
Concentration	1 mg/mL
Purity	> 95% by SDS-PAGE
Endotoxin Level	< 1.0 EU per 1 microgram of protein (determined by LAL method)
Activity	Measured in a cell proliferation assay using CTLL2 mouse cytotoxic T cells. The ED_{50} for this effect i s < 1 ng/mL.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Quality Control Testing	Loading 3 ug protein in 15% SDS-PAGE
Storage Buffer	In PBS, pH 7.4 (10% glycerol).
Storage Instruction	Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Functional Study
- SDS-PAGE

Gene Info — IL15

Entrez GeneID	3600
Protein Accession#	NP_751914
Gene Name	IL15
Gene Alias	IL-15, MGC9721
Gene Description	interleukin 15
Omim ID	600554
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Two alternatively spliced transcript variants of this gene encoding the same protein have been reported. [provided by RefSeq
Other Designations	OTTHUMP00000164617

Pathway

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Cytokine-cytokine receptor interaction
- Jak-STAT signaling pathway

Disease

- Adenocarcinoma
- Arthritis
- Asthma
- Bronchial Hyperreactivity
- Bronchiolitis
- Carcinoma
- Cardiovascular Diseases
- Celiac Disease
- Connective Tissue Diseases
- Coronary Disease
- Diabetes Mellitus
- Edema
- Esophageal Neoplasms
- Fetal Diseases
- Fractures
- Genetic Predisposition to Disease
- Hematologic Diseases
- Hodgkin Disease
- Hypersensitivity
- Hypertrophy
- Infant
- Infection

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Inflammation
- Kidney Failure
- Lung Neoplasms
- Lymphoma
- Lymphoproliferative Disorders
- Metabolic Syndrome X
- Multiple Myeloma
- Musculoskeletal Diseases
- Neoplasm
- Obesity
- Occupational Diseases
- Osteoporosis
- Pregnancy Complications
- Premature Birth
- Psoriasis
- Pulmonary Disease
- Recurrence
- Respiratory Syncytial Virus Infections
- Skin Diseases
- Urinary Bladder Neoplasms
- Waldenstrom Macroglobulinemia
- Werner syndrome

Copyright © 2023 Abnova Corporation. All Rights Reserved.