Bioactive

IL15RA (Human) Recombinant Protein

Catalog # P8350 Size 2 x 10 ug

Specification

Product Description	Human IL15RA (Q13261, 31 a.a. - 205 a.a.) partial recombinant protein with hIgG-His tag at C-terminus expressed in Sf9 cells.
Sequence	ADPITCPPPMSVEHADIWVKSYSLYSRERYICNSGFKRKAGTSSLTECVLNKATNVAHWTTPSLK CIRDPALVHQRPAPPSTVTTAGVTPQPESLSPSGKEPAASSPSSNNTAATTAAIVPGSQLMPSKS PSTGTTEISSHESSHGTPSQTTAKNWELTASASHQPPGVYPQGHSDTTLEPKSCDKTHTCPPCP APELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELT KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGKHHHHHH
Host	insect
Theoretical MW (kDa)	45.6
Form	Liquid
Preparation Method	Sf9 cell expression system
Purity	> 90% by SDS-PAGE
Activity	ED_{50} is < 2 ng/mL with Human IL-15 as determined by the ability to inhibit proliferation using CTLL2 mouse cytotoxic T cells
Recommend Usage	Biological Activity SDS-PAGE The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS pH 7.4 (10% glycerol)
Storage Instruction	Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Functional Study
- SDS-PAGE

Gene Info — IL15RA

Entrez GeneID	3601
Protein Accession#	Q13261
Gene Name	IL15RA
Gene Alias	MGC104179
Gene Description	interleukin 15 receptor, alpha
Omim ID	601070
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a cytokine receptor that specifically binds IL15 with high affinity. The receptors of IL15 and IL2 share two subunits, the IL2R beta and IL2R gamma chains. This forms the basis of many overlapping biological activities of IL15 and IL2. The IL2 receptor requires an additional IL2-specific alpha subunit for high affinity IL2 binding. This protein is structurally related to IL2R alpha, but is capable of binding IL15 with high affinity independent of other subunits, which suggests the distinct roles between IL15 and IL2. This receptor is reported to enhance cell proliferation and expression of apoptosis inhibitor BCL2L1/BCL2-XL and BCL2. Multiple alternatively spliced transcript variants of this gene have been reported. The full length sequences of only two variants encoding distinct isoforms are available. [provided by RefSeq
Other Designations	OTTHUMP00000019021|OTTHUMP00000019022

Pathway

- Cytokine-cytokine receptor interaction
- Jak-STAT signaling pathway

Disease

- Adenocarcinoma
- Alzheimer Disease

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Asthma
- Bronchiolitis
- Carcinoma
- Celiac Disease
- Esophageal Neoplasms
- Genetic Predisposition to Disease
- Hyperparathyroidism
- Infant
- Kidney Failure
- Lung Neoplasms
- Lymphoma
- Metabolic Syndrome X
- Obesity
- Pulmonary Disease
- Respiratory Syncytial Virus Infections
- Syndrome
- Urinary Bladder Neoplasms
- Werner syndrome

Copyright © 2023 Abnova Corporation. All Rights Reserved.