


# STK3 & CASP3 Protein Protein Interaction Antibody Pair

Catalog # DI0475      Size 1 Set

## Applications



Representative image of Proximity Ligation Assay of protein-protein interactions between STK3 and CASP3. HeLa cells were stained with anti-STK3 rabbit purified polyclonal antibody 1:1200 and anti-CASP3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

## Specification

<b>Product Description</b>	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the STK3 protein, and the other against the CASP3 protein for use in <a href="#">in situ Proximity Ligation Assay</a> . <a href="#">See Publication Reference below</a> .
<b>Reactivity</b>	Human
<b>Quality Control Testing</b>	Protein protein interaction immunofluorescence result. Representative image of Proximity Ligation Assay of protein-protein interactions between STK3 and CASP3. HeLa cells were stained with anti-STK3 rabbit purified polyclonal antibody 1:1200 and anti-CASP3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
<b>Supplied Product</b>	Antibody pair set content: 1. STK3 rabbit purified polyclonal antibody (100 ug) 2. CASP3 mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
<b>Storage Instruction</b>	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

## Applications

- *In situ* Proximity Ligation Assay (Cell)

## Gene Info — CASP3

Entrez GeneID	<a href="#">836</a>
Gene Name	CASP3
Gene Alias	CPP32, CPP32B, SCA-1
Gene Description	caspase 3, apoptosis-related cysteine peptidase
Omim ID	<a href="#">600636</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	<p>This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein. [provided by RefSeq]</p>
Other Designations	OTTHUMP00000165054 PARP cleavage protease SREBP cleavage activity 1 Yama apopain caspase 3 caspase 3, apoptosis-related cysteine protease cysteine protease CPP32 procaspase3

## Gene Info — STK3

Entrez GeneID	<a href="#">6788</a>
Gene Name	STK3
Gene Alias	FLJ90748, KRS1, MST2
Gene Description	serine/threonine kinase 3 (STE20 homolog, yeast)
Omim ID	<a href="#">605030</a>
Gene Ontology	<a href="#">Hyperlink</a>

**Gene Summary**

Protein kinase activation is a frequent response of cells to treatment with growth factors, chemicals, heat shock, or apoptosis-inducing agents. This protein kinase activation presumably allows cells to resist unfavorable environmental conditions. The yeast 'sterile 20' (Ste20) kinase acts upstream of the mitogen-activated protein kinase (MAPK) cascade that is activated under a variety of stress conditions. MST2 was identified as a kinase that is activated by the proapoptotic agents staurosporine and FAS ligand (MIM 134638) (Taylor et al., 1996 [PubMed 8816758]; Lee et al., 2001 [PubMed 11278283]).[supplied by OMIM]

**Other Designations**

serine/threonine kinase 3 (Ste20, yeast homolog)

## Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Colorectal cancer](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [MAPK signaling pathway](#)
- [MAPK signaling pathway](#)
- [Natural killer cell mediated cytotoxicity](#)
- [p53 signaling pathway](#)
- [Pathways in cancer](#)

## Disease

- [Adenocarcinoma](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Clubfoot](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)

- [Diabetic Nephropathies](#)
- [Disease Progression](#)
- [Edema](#)
- [Endometrial Neoplasms](#)
- [Esophageal Neoplasms](#)
- [Gastrointestinal Stromal Tumors](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Lung carcinoma](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Multiple Sclerosis](#)
- [NARP](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Occupational Diseases](#)

- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Small Cell Lung Carcinoma](#)
- [Stomach Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)