


MAP3K5 & DAXX Protein Protein Interaction Antibody Pair

Catalog # DI0525

Size 1 Set

Applications



Representative image of Proximity Ligation Assay of protein-protein interactions between MAP3K5 and DAXX. HeLa cells were stained with anti-MAP3K5 rabbit purified polyclonal antibody 1:1200 and anti-DAXX mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Specification

Product Description

This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the MAP3K5 protein, and the other against the DAXX protein for use in [in situ Proximity Ligation Assay](#). [See Publication Reference below](#).

Reactivity

Human

Quality Control Testing

Protein protein interaction immunofluorescence result.
Representative image of Proximity Ligation Assay of protein-protein interactions between MAP3K5 and DAXX. HeLa cells were stained with anti-MAP3K5 rabbit purified polyclonal antibody 1:1200 and anti-DAXX mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product

Antibody pair set content:
1. MAP3K5 rabbit purified polyclonal antibody (100 ug)
2. DAXX mouse monoclonal antibody (40 ug)
*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze-thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- *In situ* Proximity Ligation Assay (Cell)

Gene Info — DAXX

Entrez GeneID	1616
Gene Name	DAXX
Gene Alias	BING2, DAP6, EAP1, MGC126245, MGC126246
Gene Description	death-domain associated protein
Omim ID	603186
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a multifunctional protein that resides in multiple locations in the nucleus and in the cytoplasm. It interacts with a wide variety of proteins, such as apoptosis antigen Fas, centromere protein C, and transcription factor erythroblastosis virus E26 oncogene homolog 1. In the nucleus, the encoded protein functions as a potent transcription repressor that binds to sumoylated transcription factors. Its repression can be relieved by the sequestration of this protein into promyelocytic leukemia nuclear bodies or nucleoli. This protein also associates with centromeres in G2 phase. In the cytoplasm, the encoded protein may function to regulate apoptosis. The subcellular localization and function of this protein are modulated by post-translational modifications, including sumoylation, phosphorylation and polyubiquitination. Alternative splicing results in multiple transcript variants. [provided by RefSeq]
Other Designations	CENP-C binding protein ETS1-associated protein 1 Fas-binding protein OTTHUMP00000029289 OTTHUMP00000029290 death-associated protein 6

Gene Info — MAP3K5

Entrez GeneID	4217
Gene Name	MAP3K5
Gene Alias	ASK1, MAPKKK5, MEKK5
Gene Description	mitogen-activated protein kinase kinase kinase 5
Omim ID	602448
Gene Ontology	Hyperlink

Gene Summary

Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK. [provided by RefSeq]

Other Designations

MAP/ERK kinase kinase 5|MAPK/ERK kinase kinase 5|OTTHUMP00000017275|apoptosis signal regulating kinase

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [MAPK signaling pathway](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)

Disease

- [Asthma](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Diabetes Mellitus](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Hypersensitivity](#)
- [Inflammation](#)

- [Insulin Resistance](#)
- [Lupus Erythematosus](#)
- [Lymphoma](#)
- [Tobacco Use Disorder](#)