


PDGFRB & FLT1 Protein Protein Interaction Antibody Pair

Catalog # DI0624


Size 1 Set

Applications




In situ Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between PDGFRB and FLT1. PC-3 cells were stained with anti-PDGFRB rabbit purified polyclonal antibody 1:100 and anti-FLT1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).



In situ Proximity Ligation Assay (Cell)

Confocal microscopy image of Proximity Ligation Assay of protein-protein interactions between PDGFRB and FLT1. PC-3 cells were stained with anti-PDGFRB rabbit purified polyclonal antibody 1:100 and anti-FLT1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).



Representative image of Proximity Ligation Analysis of protein-protein interactions between PDGFRB and FLT1. HeLa cells were stained with anti-PDGFRB rabbit purified polyclonal antibody 1:100 and anti-FLT1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Specification

Product Description	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the PDGFRB protein, and the other against the FLT1 protein for use in in situ Proximity Ligation Assay . See Publication Reference below .
Reactivity	Human
Quality Control Testing	Protein protein interaction immunofluorescence result. Representative image of Proximity Ligation Analysis of protein-protein interactions between PDGFRB and FLT1. HeLa cells were stained with anti-PDGFRB rabbit purified polyclonal antibody 1:100 and anti-FLT1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) downloaded from The Centre for Image Analysis at Uppsala University.
Supplied Product	Antibody pair set content: 1. PDGFRB rabbit purified polyclonal antibody (100 ug) 2. FLT1 mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze-thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- *In situ* Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between PDGFRB and FLT1. PC-3 cells were stained with anti-PDGFRB rabbit purified polyclonal antibody 1:100 and anti-FLT1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

- *In situ* Proximity Ligation Assay (Cell)

Confocal microscopy image of Proximity Ligation Assay of protein-protein interactions between PDGFRB and FLT1. PC-3 cells were stained with anti-PDGFRB rabbit purified polyclonal antibody 1:100 and anti-FLT1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

Gene Info — FLT1

Entrez GeneID	2321
Gene Name	FLT1
Gene Alias	FLT, VEGFR1
Gene Description	fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor)
Omim ID	165070

Gene Ontology

[Hyperlink](#)

Gene Summary

This gene encodes a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (Ig)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia

Other Designations

fms-related tyrosine kinase 1|soluble VEGF receptor 1-14|soluble VEGFR1 variant 2|soluble VEGFR1 variant 21|vascular endothelial growth factor/vascular permeability factor receptor

Gene Info — PDGFRB

Entrez GeneID

[5159](#)

Gene Name

PDGFRB

Gene Alias

CD140B, JTK12, PDGF-R-beta, PDGFR, PDGFR1

Gene Description

platelet-derived growth factor receptor, beta polypeptide

Omim ID

[131440 173410](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. This gene is flanked on chromosome 5 by the genes for granulocyte-macrophage colony-stimulating factor and macrophage-colony stimulating factor receptor; all three genes may be implicated in the 5-q syndrome. A translocation between chromosomes 5 and 12, that fuses this gene to that of the translocation, ETV6, leukemia gene, results in chronic myeloproliferative disorder with eosinophilia. [provided by RefSeq]

Other Designations

beta platelet-derived growth factor receptor|platelet-derived growth factor receptor beta|soluble PDGFRb variant 1

Pathway

- [Calcium signaling pathway](#)
- [Colorectal cancer](#)

- [Cytokine-cytokine receptor interaction](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Focal adhesion](#)
- [Focal adhesion](#)
- [Gap junction](#)
- [Glioma](#)
- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Abortion](#)
- [Acute Disease](#)
- [Adenocarcinoma](#)
- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Cardiovascular Diseases](#)
- [Cardiovascular Diseases](#)
- [Cell Transformation](#)
- [Chorioamnionitis](#)

- [Colorectal Neoplasms](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Diabetes Mellitus](#)
- [Disease Models](#)
- [Edema](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Esophageal Neoplasms](#)
- [Fetal Growth Retardation](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Hypercholesterolemia](#)
- [Hyperparathyroidism](#)
- [Hypersensitivity](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Lymphoma](#)
- [Malaria](#)
- [Melanoma](#)
- [Metabolic Syndrome X](#)
- [Neoplasms](#)
- [Neovascularization](#)

- [Obstetric Labor](#)
- [Osteoporosis](#)
- [Ovarian Neoplasms](#)
- [Placenta Diseases](#)
- [Pre-Eclampsia](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Sarcoidosis](#)
- [Schizophrenia](#)
- [Scleroderma](#)
- [Skin Neoplasms](#)
- [Subdural Effusion](#)
- [Vaginosis](#)