
PLCG1(phospho Y783) & PLCG1 Protein Phosphorylation Antibody Pair

Catalog # DP0166 Size 1 Set

Applications



Representative image of Proximity Ligation Assay of protein phosphorylation. HeLa cells were stained with dual recognition antibody pair set, rabbit polyclonal antibody 1:1200 and mouse monoclonal antibody 1:50. Each red dot represents one single phosphorylated protein. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Specification

Product Description	This protein phosphorylation antibody pair set comes with two antibodies, one against the PLCG1 protein, and the other against the specific Y783 phosphorylated site of PLCG1 for use in *in situ* Proximity Ligation Assay. See Publication Reference below.
Reactivity	Human
Quality Control Testing	Dual recognition immunofluorescence result. Representative image of Proximity Ligation Assay of protein phosphorylation. HeLa cells were stained with dual recognition antibody pair set, rabbit polyclonal antibody 1:1200 and mouse monoclonal antibody 1:50. Each red dot represents one single phosphorylated protein. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
Supplied Products	Antibody pair set content: 1. Phospho-PLCG1 Y783 rabbit polyclonal antibody (20 ul) In PBS, 150 mM NaCl, pH 7.4 (0.02% sodium azide, 50% glycerol) 2. PLCG1 mouse monoclonal antibody (40 ug) In 1x PBS, pH 7.2 *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Applications

- *In situ* Proximity Ligation Assay (Cell)

Gene Info — PLCG1

Entrez GeneID	5335
Gene Name	PLCG1
Gene Alias	PLC-II, PLC1, PLC148, PLCgamma1
Gene Description	phospholipase C, gamma 1
Omim ID	172420
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq
Other Designations	1-phosphatidyl-D-myo-inositol-4,5-bisphosphate|1-phosphatidylinositol-4,5-bisphosphate phosphodiesterase gamma 1|OTTHUMP00000031787|OTTHUMP00000178982|PLC-gamma-1|inositol trisphosphohydrolase|monophosphatidylinositol phosphodiesterase|phosphatidylinositol

Pathway

- Calcium signaling pathway
- Epithelial cell signaling in Helicobacter pylori infection
- ErbB signaling pathway
- Fc epsilon RI signaling pathway
- Fc gamma R-mediated phagocytosis
- Glioma
- Inositol phosphate metabolism

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Leukocyte transendothelial migration
- Metabolic pathways
- Natural killer cell mediated cytotoxicity
- Neurotrophin signaling pathway
- Non-small cell lung cancer
- Pathways in cancer
- Phosphatidylinositol signaling system
- T cell receptor signaling pathway
- VEGF signaling pathway
- Vibrio cholerae infection

Disease

- Bipolar Disorder
- Cardiovascular Diseases
- Diabetes Mellitus
- Edema
- Genetic Predisposition to Disease
- HIV Infections
- Kidney Failure
- Mental Disorders
- Multiple Sclerosis

Copyright © 2023 Abnova Corporation. All Rights Reserved.