


ACVR1B monoclonal antibody (M06), clone 2E12

Catalog # H00000091-M06

Size 100 ug

Applications




Western Blot (Transfected lysate)

Western Blot analysis of ACVR1B expression in transfected 293T cell line by ACVR1B monoclonal antibody (M06), clone 2E12.


Lane 1: ACVR1B transfected lysate (Predicted MW: 56.8 kDa).

Lane 2: Non-transfected lysate.



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged ACVR1B is approximately 0.03ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.96 kDa) .

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant ACVR1B.

Immunogen	ACVR1B (AAH00254, 24 a.a. ~ 126 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	SGPRGVQALLCACTSCLQANYTCETDGACMVSIFNLDGMEHHVRTCIPKVELVPAGKPFYCLSS EDLRNTHCCYTDYCNRIDLRVPSGHLKEPEHPSMWGPVE
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (93); Rat (95)
Isotype	IgG1 Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.96 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of ACVR1B expression in transfected 293T cell line by ACVR1B monoclonal antibody (M06), clone 2E12.

Lane 1: ACVR1B transfected lysate (Predicted MW: 56.8 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged ACVR1B is approximately 0.03ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — ACVR1B

Entrez GeneID	91
GeneBank Accession#	BC000254
Protein Accession#	AAH00254
Gene Name	ACVR1B
Gene Alias	ACTRIB, ACVRLK4, ALK4, SKR2
Gene Description	activin A receptor, type IB
Omim ID	601300
Gene Ontology	Hyperlink
Gene Summary	<p>Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with a cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling, and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. This gene encodes activin A type IB receptor, composed of 11 exons. Alternative splicing and alternative polyadenylation result in 3 fully described transcript variants. The mRNA expression of variants 1, 2, and 3 is confirmed, and a potential fourth variant contains an alternative exon 8 and lacks exons 9 through 11, but its mRNA expression has not been confirmed. [provided by RefSeq]</p>
Other Designations	activin A receptor, type II-like kinase 4 activin A type IB receptor activin receptor-like kinase 4 serine(threonine) protein kinase receptor R2

Pathway

- [Adherens junction](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [MAPK signaling pathway](#)

- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [TGF-beta signaling pathway](#)

Disease

- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Obesity](#)
- [Ovarian Failure](#)
- [Polycystic Ovary Syndrome](#)
- [Puberty](#)
- [Schizophrenia](#)
- [Thrombophilia](#)
- [Tobacco Use Disorder](#)