

ADH4 rabbit monoclonal antibody

Catalog # H00000127-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human ADH4 peptide using ARM Technology.
Immunogen	A synthetic peptide of human ADH4 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human ADH4 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — ADH4

Entrez GeneID	127
GeneBank Accession#	ADH4
Gene Name	ADH4
Gene Alias	ADH-2
Gene Description	alcohol dehydrogenase 4 (class II), pi polypeptide
Omim ID	103740
Gene Ontology	Hyperlink
Gene Summary	This gene encodes class II alcohol dehydrogenase 4 pi subunit, which is a member of the alcohol dehydrogenase family. Members of this enzyme family metabolize a wide variety of substrates, including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products. Class II alcohol dehydrogenase is a homodimer composed of 2 pi subunits. It exhibits a high activity for oxidation of long-chain aliphatic alcohols and aromatic alcohols and is less sensitive to pyrazole. This gene is localized to chromosome 4 in the cluster of alcohol dehydrogenase genes. [provided by RefSeq]
Other Designations	aldehyde reductase class II alcohol dehydrogenase 4, pi subunit

Pathway

- [1- and 2-Methylnaphthalene degradation](#)
- [3-Chloroacrylic acid degradation](#)
- [Drug metabolism - cytochrome P450](#)
- [Fatty acid metabolism](#)
- [Glycolysis / Gluconeogenesis](#)
- [Metabolic pathways](#)
- [Metabolism of xenobiotics by cytochrome P450](#)
- [Retinol metabolism](#)

- [Tyrosine metabolism](#)

Disease

- [Adenocarcinoma](#)
- [Alcohol-Induced Disorders](#)
- [Alcoholism](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Cluster Headache](#)
- [Disease Models](#)
- [Diseases in Twins](#)
- [Esophageal Neoplasms](#)
- [Extraversion \(Psychology\)](#)
- [Flushing](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hearing Loss](#)
- [Kidney Failure](#)
- [Parkinson disease](#)
- [Personality Inventory](#)
- [Substance-Related Disorders](#)