

ATF4 rabbit monoclonal antibody

Catalog # H00000468-K Size 100 ug x up to 3

Specification

| | |
|--------------------------------|--|
| Product Description | Rabbit monoclonal antibody raised against a human ATF4 peptide using ARM Technology. |
| Immunogen | A synthetic peptide of human ATF4 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence. |
| Host | Rabbit |
| Library Construction | Non-fusion antibody library from rabbit spleen (ARM Technology). |
| Expression | Overexpression vector and transfection into 293H cell line. |
| Reactivity | Human |
| Purification | Protein A |
| Isotype | IgG |
| Quality Control Testing | Antibody reactive against human ATF4 peptide by ELISA and mammalian transfected lysate by Western Blot. |
| Storage Buffer | In 1x PBS, pH 7.4 |
| Storage Instruction | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. |
| Deliverable | Up to three rabbit IgG clones of 100 ug each will be delivered to customer. |
| Note | 1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request. |

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — ATF4

Entrez GeneID [468](#)

GeneBank Accession# [ATF4](#)

Gene Name ATF4

Gene Alias CREB-2, CREB2, TAXREB67, TXREB

Gene Description activating transcription factor 4 (tax-responsive enhancer element B67)

Omim ID [604064](#)

Gene Ontology [Hyperlink](#)

Gene Summary

This gene encodes a transcription factor that was originally identified as a widely expressed mammalian DNA binding protein that could bind a tax-responsive enhancer element in the LTR of HTLV-1. The encoded protein was also isolated and characterized as the cAMP-response element binding protein 2 (CREB-2). The protein encoded by this gene belongs to a family of DNA-binding proteins that includes the AP-1 family of transcription factors, cAMP-response element binding proteins (CREBs) and CREB-like proteins. These transcription factors share a leucine zipper region that is involved in protein-protein interactions, located C-terminal to a stretch of basic amino acids that functions as a DNA binding domain. Two alternative transcripts encoding the same protein have been described. Two pseudogenes are located on the X chromosome at q28 in a region containing a large inverted duplication. [provided by RefSeq]

Other Designations activating transcription factor 4|cAMP response element-binding protein 2

Pathway

- [GnRH signaling pathway](#)
- [Long-term potentiation](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Prostate cancer](#)

Disease

- [Bipolar Disorder](#)
- [Genetic Predisposition to Disease](#)
- [Mental Disorders](#)
- [Neuropsychological Tests](#)
- [Schizophrenia](#)