


BDH1 monoclonal antibody (M03A), clone 4B3


Catalog # H00000622-M03A Size 200 uL

Applications



Western Blot (Cell lysate)

BDH1 monoclonal antibody (M03A), clone 4B3. Western Blot analysis of BDH1 expression in Raw 264.7 (Cat # L024V1).



Western Blot detection against Immunogen (63.47 KDa).

Specification	
Product Description	Mouse monoclonal antibody raised against a full-length recombinant BDH1.
Immunogen	BDH1 (AAH11964, 1 a.a. ~ 343 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	MLATRLSRPLSRLPGKTLSACDRENGARRPLLLGSTSFIPIGRRTYASAAEPVGSKAVLVTGCDSG FGFSLAKHLHSKGFLVFAGCLMKDKGHDGVKELDSLNSDRLRTVQLNVCSSEEVEKVVEIVRSS LKDPEKGMWGLVNNAGISTFGEVEFTSLETYKQVAEVNLWGTVRMTKSFLPLIRRAKGRVVNISS MLGRMANPARSPYCITKFGVEAFSDCLRYEMYPLGVKVSVVEPGNFIAATSLYSPESIQAIAKKMW EELPEVVRKDYGKKYFDEKIAKMETYCSSGSTDTSPVIDAVTHALTATTPYTRYHPMDYYWWLRM QIMTHLPGAISDMIYIR
Host	Mouse



Product Information

Reactivity	Human, Mouse
Interspecies Antigen Sequence	Mouse (86); Rat (87)
Isotype	lgG2a Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (63.47 KDa).
Storage Buffer	In ascites fluid
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

Western Blot (Cell lysate)

 $BDH1\ monoclonal\ antibody\ (M03A),\ clone\ 4B3.\ Western\ Blot\ analysis\ of\ BDH1\ expression\ in\ Raw\ 264.7\ (\ Cat\ \#\ L024V1\).$

Protocol Download

Western Blot (Recombinant protein)

Protocol Download

ELISA

Gene Info — BDH1	
Entrez GeneID	<u>622</u>
GeneBank Accession#	BC011964
Protein Accession#	AAH11964
Gene Name	BDH1
Gene Alias	BDH, MGC2723, MGC4347, MGC9788, SDR9C1
Gene Description	3-hydroxybutyrate dehydrogenase, type 1
Omim ID	603063
Gene Ontology	<u>Hyperlink</u>



Product Information

Gene Summary

This gene encodes a member of the short-chain dehydrogenase/reductase gene family. The encoded protein forms a homotetrameric lipid-requiring enzyme of the mitochondrial membrane and has a specific requirement for phosphatidylcholine for optimal enzymatic activity. The encoded protein catalyzes the interconversion of acetoacetate and (R)-3-hydroxybutyrate, the two major ketone bodies produced during fatty acid catabolism. Alternatively spliced transcript variants encoding the same protein have been described. [provided by RefSeq

Other Designations

(R)-3-hydroxybutyrate dehydrogenase|3-hydroxybutyrate dehydrogenase|3-hydroxybutyrate dehydrogenase|3-hydroxybutyrate dehydrogenase, mitochondrial|short chain dehydrogenase/reductase family 9C, member 1

Pathway

- Butanoate metabolism
- Metabolic pathways
- Synthesis and degradation of ketone bodies