

## CD8B rabbit monoclonal antibody

Catalog # H00000926-K

Size 100 ug x up to 3

### Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against a human CD8B peptide using ARM Technology.
<b>Immunogen</b>	A synthetic peptide of human CD8B is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
<b>Host</b>	Rabbit
<b>Library Construction</b>	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
<b>Expression</b>	Overexpression vector and transfection into 293H cell line.
<b>Reactivity</b>	Human
<b>Purification</b>	Protein A
<b>Isotype</b>	IgG
<b>Quality Control Testing</b>	Antibody reactive against human CD8B peptide by ELISA and mammalian transfected lysate by Western Blot.
<b>Storage Buffer</b>	In 1x PBS, pH 7.4
<b>Storage Instruction</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Deliverable</b>	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
<b>Note</b>	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

### Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — CD8B

Entrez GeneID [926](#)

GeneBank Accession# [CD8B](#)

Gene Name CD8B

Gene Alias CD8B1, LYT3, Leu2, Ly3, MGC119115

Gene Description CD8b molecule

Omim ID [186730](#)

Gene Ontology [Hyperlink](#)

**Gene Summary**

The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T-cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 beta chain isoforms. Multiple alternatively spliced transcript variants encoding distinct membrane associated or secreted isoforms have been described. A pseudogene, also located on chromosome 2, has been identified. [provided by RefSeq]

**Other Designations**

CD8 antigen, beta polypeptide (p37)|CD8 antigen, beta polypeptide 1 (p37)|CD8b antigen|OTTHUMP00000160761|T lymphocyte surface glycoprotein beta chain|T-cell surface glycoprotein CD8 beta chain

## Pathway

- [Antigen processing and presentation](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Hematopoietic cell lineage](#)
- [Primary immunodeficiency](#)
- [T cell receptor signaling pathway](#)

## Disease

- [Tobacco Use Disorder](#)