


CD86 monoclonal antibody (M01), clone 3D7

Catalog # H00000942-M01

Size 100 ug

Applications



Immunoprecipitation

Immunoprecipitation of CD86 transfected lysate using anti-CD86 monoclonal antibody and Protein A Magnetic Bead, and immunoblotted with CD86 MaxPab rabbit polyclonal antibody.

Specification

Product Description	Mouse monoclonal antibody raised against a partial recombinant CD86.
Immunogen	CD86 (AAH40261, 220 a.a. ~ 329 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	LETDKTRLLSSPFSIELEDPPPPDHIPWITAVLPTVIICVMVFCLILWKWKKKRPRNSYKCGTNT MEREESEQTKKREKIHIPERSDETQRVFKSSKTSSCDKSDTCF
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (36); Rat (31)
Isotype	IgG2a Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Immunoprecipitation

Immunoprecipitation of CD86 transfected lysate using anti-CD86 monoclonal antibody and Protein A Magnetic Bead, and immunoblotted with CD86 MaxPab rabbit polyclonal antibody.

[Protocol Download](#)

- ELISA

Gene Info — CD86

Entrez GeneID [942](#)

GeneBank Accession# [BC040261](#)

Protein Accession# [AAH40261](#)

Gene Name CD86

Gene Alias B7-2, B70, CD28LG2, LAB72, MGC34413

Gene Description CD86 molecule

Omim ID [601020](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in two transcript variants encoding different isoforms. Additional transcript variants have been described, but their full-length sequences have not been determined. [provided by RefSeq]

Other Designations B-lymphocyte activation antigen B7-2|B7-2 antigen|CD28 antigen ligand 2|CD86 antigen|CD86 antigen (CD28 antigen ligand 2, B7-2 antigen)|CTLA-4 counter-receptor B7.2

Pathway

- [Allograft rejection](#)

- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)
- [Type I diabetes mellitus](#)

Disease

- [Acute Disease](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Colorectal Neoplasms](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Drug Eruptions](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [Hypersensitivity](#)

- [Infant](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Multiple Sclerosis](#)
- [Papillomavirus Infections](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Scleroderma](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Werner syndrome](#)