


CD86 (Human) Recombinant Protein (Q01)

Catalog # H00000942-Q01

Size 25 ug, 10 ug

Applications



Specification

Product Description	Human CD86 partial ORF (AAH40261, 220 a.a. - 329 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	LETDKTRLLSSPFSIELEDPPPPDHIPWITAVLPTVIICVMVFCLILWKWKKKKRPRNSYKCGTNT MEREESEQTKKREKIHIPERSDETQRVFKSSKTSSCDKSDTCF
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	37.51
Interspecies Antigen Sequence	Mouse (36); Rat (27)
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — CD86

Entrez GeneID [942](#)

GeneBank Accession# [BC040261](#)

Protein Accession# [AAH40261](#)

Gene Name CD86

Gene Alias B7-2, B70, CD28LG2, LAB72, MGC34413

Gene Description CD86 molecule

Omim ID [601020](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in two transcript variants encoding different isoforms. Additional transcript variants have been described, but their full-length sequence has not been determined. [provided by RefSeq]

Other Designations B-lymphocyte activation antigen B7-2|B7-2 antigen|CD28 antigen ligand 2|CD86 antigen|CD86 antigen (CD28 antigen ligand 2, B7-2 antigen)|CTLA-4 counter-receptor B7.2

Pathway

- [Allograft rejection](#)

- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)
- [Type I diabetes mellitus](#)

Disease

- [Acute Disease](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Colorectal Neoplasms](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Drug Eruptions](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [Hypersensitivity](#)
- [Infant](#)

- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Multiple Sclerosis](#)
- [Papillomavirus Infections](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Scleroderma](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Werner syndrome](#)