


CD40 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00000958-T01


Size 100 uL

Applications



SDS-PAGE Gel

CD40 transfected lysate



Western Blot

Lane 1: CD40 transfected lysate (30.47 KDa).

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-CD40 full-length
Host	Human
Theoretical MW (kDa)	30.47
Interspecies Antigen Sequence	Mouse (61); Rat (59)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-CD40 antibody ([H00000958-B01](#)) by Western Blots.
SDS-PAGE Gel
CD40 transfected lysate
Western Blot
Lane 1: CD40 transfected lysate (30.47 KDa).
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — CD40

Entrez GeneID

[958](#)

GeneBank Accession#

[NM_001250](#)

Protein Accession#

[NP_001241](#)

Gene Name

CD40

Gene Alias

Bp50, CDW40, MGC9013, TNFRSF5, p50

Gene Description

CD40 molecule, TNF receptor superfamily member 5

Omim ID

[109535 606843](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been found to be essential in mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]

Other Designations

B cell surface antigen CD40|B cell-associated molecule|CD40 antigen|CD40 antigen (TNF receptor superfamily member 5)|CD40 type II isoform|CD40L receptor|OTTHUMP00000031699|OTTHUMP00000031700|nerve growth factor receptor-related B-lymphocyte activation mol

Pathway

- [Allograft rejection](#)
- [Asthma](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Cytokine-cytokine receptor interaction](#)
- [Primary immunodeficiency](#)
- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Autoimmune Diseases](#)
- [Brain Ischemia](#)
- [Bronchiolitis](#)
- [Calcinosis](#)
- [Cardiovascular Diseases](#)
- [Colitis](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Crohn Disease](#)

- [Diabetes Mellitus](#)
- [Diabetic Angiopathies](#)
- [Disease Progression](#)
- [Drug Eruptions](#)
- [Edema](#)
- [Endometriosis](#)
- [Genetic Predisposition to Disease](#)
- [Graves Disease](#)
- [Hashimoto Disease](#)
- [Hematologic Diseases](#)
- [Hepatitis B](#)
- [Hodgkin Disease](#)
- [Hypertension](#)
- [Infant](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Multiple Sclerosis](#)
- [Myasthenia Gravis](#)
- [Occupational Diseases](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)

- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Rupture](#)
- [Stroke](#)
- [Thyroiditis](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Wegener Granulomatosis](#)
- [Werner syndrome](#)