

CEBPB rabbit monoclonal antibody

Catalog # H00001051-K Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human CEBPB peptide using ARM Technology.
Immunogen	A synthetic peptide of human CEBPB is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (<u>ARM Technology</u>).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human CEBPB peptide by ELISA and mammalian transfected lysate by W estern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — CEBPB	
Entrez GenelD	1051
GeneBank Accession#	<u>CEBPB</u>
Gene Name	CEBPB
Gene Alias	C/EBP-beta, CRP2, IL6DBP, LAP, MGC32080, NF-IL6, TCF5
Gene Description	CCAAT/enhancer binding protein (C/EBP), beta
Omim ID	189965
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this intronless gene is a bZIP transcription factor which can bind as a ho modimer to certain DNA regulatory regions. It can also form heterodimers with the related protein s CEBP-alpha, CEBP-delta, and CEBP-gamma. The encoded protein is important in the regulati on of genes involved in immune and inflammatory responses and has been shown to bind to the IL -1 response element in the IL-6 gene, as well as to regulatory regions of several acute-phase and cytokine genes. In addition, the encoded protein can bind the promoter and upstream element an d stimulate the expression of the collagen type I gene. [provided by RefSeq
Other Designations	CCAAT/enhancer binding protein beta interleukin 6-dependent DNA-binding protein liver-enriche d transcriptional activator protein nuclear factor of interleukin 6 transcription factor 5

Disease

- Genetic Predisposition to Disease
- Obesity