


CST3 monoclonal antibody, clone 1H4

Catalog # H00001471-M04

Size 100 ug

Applications



Western Blot detection against Immunogen (13 kDa)

Specification

Product Description	Mouse monoclonal antibody raised against a full length native CST3.
Immunogen	Native purified human CST3.
Host	Mouse
Reactivity	Human
Form	Liquid
Isotype	IgG2a, kappa
Quality Control Testing	Antibody Reactive Against Native Protein Western Blot detection against Immunogen (13 kDa)
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Recombinant protein)

[Protocol Download](#)

- ELISA

Gene Info — CST3

Entrez GeneID [1471](#)

Gene Name CST3

Gene Alias ARMD11, MGC117328

Gene Description cystatin C

Omim ID [105150 604312](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes the most abundant extracellular inhibitor of cysteine proteases, which is found in high concentrations in biological fluids and is expressed in virtually all organs of the body. A mutation in this gene has been associated with amyloid angiopathy. Expression of this protein in vascular wall smooth muscle cells is severely reduced in both atherosclerotic and aneurysmal aortic lesions, establishing its role in vascular disease. [provided by RefSeq]

Other Designations

OTTHUMP00000030440|OTTHUMP00000164181|OTTHUMP00000164182|bA218C14.4 (cystatin C)|cystatin 3|cystatin C (amyloid angiopathy and cerebral hemorrhage)|gamma-trace|neuroendocrine basic polypeptide|post-gamma-globulin

Disease

- [Albuminuria](#)
- [Alzheimer disease](#)

- [Amnesia](#)
- [Amyloidosis](#)
- [Amyotrophic lateral sclerosis](#)
- [Angina](#)
- [Atherosclerosis](#)
- [Cardiovascular Diseases](#)
- [Cognition Disorders](#)
- [Coronary Disease](#)
- [Dementia](#)
- [Diabetes Complications](#)
- [Frontotemporal Lobar Degeneration](#)
- [Genetic Predisposition to Disease](#)
- [Lewy Body Disease](#)
- [Macular Degeneration](#)
- [Mental Status Schedule](#)
- [Metabolic Syndrome X](#)
- [Mouth Neoplasms](#)
- [Myocardial Infarction](#)
- [Neoplasms](#)
- [Neuropsychological Tests](#)
- [Oral Submucous Fibrosis](#)
- [Osteoporosis](#)
- [Parkinson Disease](#)
- [Precancerous Conditions](#)
- [Psychiatric Status Rating Scales](#)