


FHIT monoclonal antibody (M03), clone 1E3

Catalog # H00002272-M03


Size 100 ug

Applications




Western Blot (Cell lysate)

FHIT monoclonal antibody (M01), clone 1E3 Western Blot analysis of FHIT expression in HL-60 (Cat # L014V1).



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged FHIT is approximately 0.03ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.63 KDa) .

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant FHIT.

Immunogen	FHIT (AAH32336, 31 a.a. ~ 130 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	VVPGHVLVCPLRPVERFHDLRPDEVADLFQTTQRVGTVVEKHFHGTSLTFSMQDGPEAGQTVK HVHVHVLPRKAGDFHRNDSIYEELQKHKEDFPASWR
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (87); Rat (85)
Isotype	IgG1 kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

FHIT monoclonal antibody (M01), clone 1E3 Western Blot analysis of FHIT expression in HL-60 (Cat # L014V1).

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged FHIT is approximately 0.03ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — FHIT

Entrez GeneID [2272](#)

GeneBank Accession#	BC032336
Protein Accession#	AAH32336
Gene Name	FHIT
Gene Alias	AP3Aase, FRA3B
Gene Description	fragile histidine triad gene
Omim ID	601153
Gene Ontology	Hyperlink
Gene Summary	This gene, a member of the histidine triad gene family, encodes a diadenosine 5',5'''-P1,P3-triphosphate hydrolase involved in purine metabolism. The gene encompasses the common fragile site FRA3B on chromosome 3, where carcinogen-induced damage can lead to translocations and aberrant transcripts of this gene. In fact, aberrant transcripts from this gene have been found in about half of all esophageal, stomach, and colon carcinomas. [provided by RefSeq]
Other Designations	AP3A hydrolase bis(5'-adenosyl)-triphosphatase diadenosine 5',5'''-P1,P3-triphosphate hydrolase dinucleosidetriphosphatase tumor suppressor protein

Pathway

- [Non-small cell lung cancer](#)
- [Purine metabolism](#)
- [Small cell lung cancer](#)

Disease

- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Cell Transformation](#)
- [Cervical Intraepithelial Neoplasia](#)
- [Chagas Disease](#)
- [Diabetes Mellitus](#)
- [Edema](#)

- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Laryngeal Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Mouth Neoplasms](#)
- [Overweight](#)
- [Papillomavirus Infections](#)
- [Pharyngeal Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Uterine Cervical Neoplasms](#)