


HLA-DMB monoclonal antibody (M09), clone 6E10

Catalog # H00003109-M09

Size 100 ug

Applications




Western Blot (Transfected lysate)

Western Blot analysis of HLA-DMB expression in transfected 293T cell line by HLA-DMB monoclonal antibody (M09), clone 6E10.


Lane 1: HLA-DMB transfected lysate(28.9 KDa).

Lane 2: Non-transfected lysate.



Immunoprecipitation

Immunoprecipitation of HLA-DMB transfected lysate using anti-HLA-DMB monoclonal antibody and Protein A Magnetic Bead, and immunoblotted with HLA-DMB MaxPab rabbit polyclonal antibody.



Western Blot detection against Immunogen (37.62 KDa) .

Specification

Product Description

Mouse monoclonal antibody raised against a partial recombinant HLA-DMB.

Immunogen	HLA-DMB (NP_002109, 22 a.a. ~ 129 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	VAHVESTCLLDDAGTPKDFTYCISFNKDLLTCWDPEENKMAPCEFGVLNSLANVLSQHLNQKDT LMQRLRNGLQNCATHTQPFWGSLTNRTRPPSVQVAKTTPTNTRE
Host	Mouse
Reactivity	Human
Isotype	IgG2b Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.62 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of HLA-DMB expression in transfected 293T cell line by HLA-DMB monoclonal antibody (M09), clone 6E10.

Lane 1: HLA-DMB transfected lysate(28.9 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Immunoprecipitation

Immunoprecipitation of HLA-DMB transfected lysate using anti-HLA-DMB monoclonal antibody and Protein A Magnetic Bead, and immunoblotted with HLA-DMB MaxPab rabbit polyclonal antibody.

[Protocol Download](#)

- ELISA

Gene Info — HLA-DMB

Entrez GeneID

[3109](#)

GeneBank Accession#	NM_002118
Protein Accession#	NP_002109
Gene Name	HLA-DMB
Gene Alias	D6S221E, RING7
Gene Description	major histocompatibility complex, class II, DM beta
Omim ID	142856
Gene Ontology	Hyperlink
Gene Summary	HLA-DMB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DMA) and a beta (DMB) chain, both anchored in the membrane. It is located in intracellular vesicles. DM plays a central role in the peptide loading of MHC class II molecules by helping to release the CLIP (class II-associated invariant chain peptide) molecule from the peptide binding site. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. [provided by RefSeq]
Other Designations	MHC class II HLA-DMB MHC class II antigen HLA-DM beta chain OTTHUMP00000029257 class II histocompatibility antigen, M beta chain

Pathway

- [Allograft rejection](#)
- [Antigen processing and presentation](#)
- [Asthma](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Systemic lupus erythematosus](#)
- [Type I diabetes mellitus](#)

Disease

- [Antiphospholipid Syndrome](#)
- [Arthritis](#)
- [Cardiovascular Diseases](#)
- [Condylomata Acuminata](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Lupus Erythematosus](#)
- [Measles](#)
- [Psoriasis](#)