


HLA-DMB (Human) IP-WB Antibody Pair

Catalog # H00003109-PW4

Size 1 Set

Applications



Immunoprecipitation of HLA-DMB transfected lysate using mouse monoclonal anti-HLA-DMB and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with rabbit polyclonal anti-HLA-DMB.

Specification

Product Description	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.
Reactivity	Human
Quality Control Testing	Immunoprecipitation-Western Blot (IP-WB) Immunoprecipitation of HLA-DMB transfected lysate using mouse monoclonal anti-HLA-DMB and Protein A Magnetic Bead (U0007), and immunoblotted with rabbit polyclonal anti-HLA-DMB.
Supplied Product	Antibody pair set content: 1. Antibody pair for IP: mouse monoclonal anti-HLA-DMB (300 ug) 2. Antibody pair for WB: rabbit polyclonal anti-HLA-DMB (50 ul)
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- Immunoprecipitation-Western Blot

[Protocol Download](#)

Gene Info — HLA-DMB

Entrez GeneID [3109](#)

Gene Name HLA-DMB

Gene Alias D6S221E, RING7

Gene Description major histocompatibility complex, class II, DM beta

Omim ID [142856](#)

Gene Ontology [Hyperlink](#)

Gene Summary

HLA-DMB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DMA) and a beta (DMB) chain, both anchored in the membrane. It is located in intracellular vesicles. DM plays a central role in the peptide loading of MHC class II molecules by helping to release the CLIP (class II-associated invariant chain peptide) molecule from the peptide binding site. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. [provided by RefSeq]

Other Designations MHC class II HLA-DMB|MHC class II antigen HLA-DM beta chain|OTTHUMP00000029257|class II histocompatibility antigen, M beta chain

Pathway

- [Allograft rejection](#)
- [Antigen processing and presentation](#)
- [Asthma](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Systemic lupus erythematosus](#)
- [Type I diabetes mellitus](#)

Disease

- [Antiphospholipid Syndrome](#)
- [Arthritis](#)
- [Cardiovascular Diseases](#)
- [Condylomata Acuminata](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Lupus Erythematosus](#)
- [Measles](#)
- [Psoriasis](#)