

DNAxPAb

Hard-to-Find
Antibody

CXCR2 DNAxPab

Catalog # H00003579-W01P

Size 200 ug

Specification

Product Description	Rabbit polyclonal antibody raised against a partial-length human CXCR2 DNA using DNAx™ Immune technology.
Technology	DNAx™ Immune
Immunogen	Extracellular membrane domain (ECD) human DNA
Host	Rabbit
Reactivity	Human
Purification	Protein A
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- Immunofluorescence (Transfected cell)
- Flow Cytometry (Transfected cell)

Gene Info — CXCR2

Entrez GeneID	3579
GeneBank Accession#	NM_001557.2
Protein Accession#	NP_001548.1
Gene Name	CXCR2
Gene Alias	CD182, CDw128b, CMKAR2, IL8R2, IL8RA, IL8RB
Gene Description	chemokine (C-X-C motif) receptor 2
Omim ID	146928
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. This gene, IL8RA, a gene encoding another high affinity IL8 receptor, as well as IL8RBP, a pseudogene of IL8RB, form a gene cluster in a region mapped to chromosome 2q33-q36. [provided by RefSeq]
Other Designations	C-X-C chemokine receptor type 2 CXC-R2 CXCR-2 CXCR2 gene for IL8 receptor type B GRO/MGSA receptor IL-8 receptor type 2 IL-8RB chemokine (CXC) receptor 2 high affinity interleukin-8 receptor B interleukin 8 receptor B interleukin 8 receptor type 2 interl

Pathway

- [Chemokine signaling pathway](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)

Disease

- [Acquired Immunodeficiency Syndrome](#)
- [Adenocarcinoma](#)

- [Alcoholism](#)
- [Asthma](#)
- [Bacteriuria](#)
- [Behcet Syndrome](#)
- [Biliary Tract Neoplasms](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Conduct Disorder](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Helicobacter Infections](#)
- [Hematologic Diseases](#)
- [Hepatitis C](#)

- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Infant](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Leishmaniasis](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Maduromycosis](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Multiple Sclerosis](#)
- [Neoplasm Invasiveness](#)
- [Obstetric Labor](#)
- [Occupational Diseases](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Papillomavirus Infections](#)
- [Periodontal Diseases](#)
- [Periodontitis](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)

- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoma](#)
- [Stomach Neoplasms](#)
- [Tuberculosis](#)
- [Uterine Cervical Neoplasms](#)
- [Viremia](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)