

NDUFA5 (Human) Matched Antibody Pair

Catalog # H00004698-AP21 Size 1 Set

Applications



Sandwich ELISA detection sensitivity ranging from 3 ng/ml to 100 ng/ml.

Specification	
Product Description	This antibody pair set comes with a matched antibody pair to detect and quantify the protein level of human NDUFA5.
Reactivity	Human
Quality Control Testing	Standard curve using recombinant protein (H00004698-P01) as an analyte. Sandwich ELISA detection sensitivity ranging from 3 ng/ml to 100 ng/ml.
Supplied Product	Antibody pair set content: 1. Capture antibody: rabbit MaxPab® affinity purified polyclonal anti-NDUFA5 (100 ug) 2. Detection antibody: mouse polyclonal anti-NDUFA5 (40 ul) *Reagents are sufficient for at least 3-5 x 96 well plates using recommended protocols.
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze tha w cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

• ELISA Pair (Recombinant protein)

Protocol Download

😵 Abnova

Product Information

Gene Info — NDUFA5

Entrez GenelD	<u>4698</u>
Gene Name	NDUFA5
Gene Alias	B13, CI-13KD-B, DKFZp781K1356, FLJ12147, NUFM, UQOR13
Gene Description	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 5, 13kDa
Omim ID	<u>601677</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The human NDUFA5 gene codes for the B13 subunit of complex I of the respiratory chain, which tr ansfers electrons from NADH to ubiquinone. The high degree of conservation of NDUFA5 extendi ng to plants and fungi indicates its functional significance in the enzyme complex. The protein loca lizes to the inner mitochondrial membrane as part of the 7 component-containing, water soluble "ir on-sulfur protein" (IP) fraction of complex I, although its specific role is unknown. It is assumed to u ndergo post-translational removal of the initiator methionine and N-acetylation of the next amino a cid. The predicted secondary structure is primarily alpha helix, but the carboxy-terminal half of the protein has high potential to adopt a coiled-coil form. The amino-terminal part contains a putative beta sheet rich in hydrophobic amino acids that may serve as mitochondrial import signal. Relate d pseudogenes have also been identified on four other chromosomes. [provided by RefSeq
Other Designations	Complex I-13KD-B NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 5 NADH dehydroge nase (ubiquinone) 1 alpha subcomplex, 5 (13kD, B13) type I dehydrogenase ubiquinone reductas e

Pathway

- Metabolic pathways
- Oxidative phosphorylation

Disease

- Genetic Predisposition to Disease
- Prostatic Neoplasms
- Tobacco Use Disorder