


ABCC8 monoclonal antibody (M02), clone 3E1

Catalog # H00006833-M02


Size 100 ug

Applications



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged ABCC8 is 0.03 ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.74 KDa) .

Specification

Product Description	Mouse monoclonal antibody raised against a partial recombinant ABCC8.
Immunogen	ABCC8 (NP_000343, 611 a.a. ~ 710 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	SEFLSSAEIREEQCAPHEPTQGPASKYQAVPLRVVNRKRPAREDCRGLTGPLQSLVPSADGDA DNCCVQIMGGYFTWTPDGIPTLSNITIRIPRGQLTM
Host	Mouse
Reactivity	Human

Interspecies Antigen Sequence	Mouse (85); Rat (85)
Isotype	IgG1 Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged ABCC8 is 0.03 ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — ABCC8

Entrez GeneID	6833
GeneBank Accession#	NM_000352
Protein Accession#	NP_000343
Gene Name	ABCC8
Gene Alias	ABC36, HHF1, HI, HRINS, MRP8, PHHI, SUR, SUR1, TNDM2
Gene Description	ATP-binding cassette, sub-family C (CFTR/MRP), member 8
Omim ID	125853 240800 256450 600509 606176 610374
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MRP subfamily which is involved in multi-drug resistance. This protein functions as a modulator of ATP-sensitive potassium channels and insulin release. Mutations and deficiencies in this protein have been observed in patients with hyperinsulinemic hypoglycemia of infancy, an autosomal recessive disorder of unregulated and high insulin secretion. Mutations have also been associated with non-insulin-dependent diabetes mellitus type II, an autosomal dominant disease of defective insulin secretion. Alternative splicing of this gene has been observed; however, the transcript variants have not been fully described. [provided by RefSeq]

Other Designations

ATP-binding cassette, sub-family C, member 8|sulfonylurea receptor (hyperinsulinemia)

Pathway

- [ABC transporters](#)
- [Type II diabetes mellitus](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Atherosclerosis](#)
- [Cardiovascular Diseases](#)
- [Cerebrovascular Accident](#)
- [Chronic Disease](#)
- [Diabetes](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Diabetic Angiopathies](#)
- [Disease Progression](#)
- [Drug Toxicity](#)
- [Edema](#)

- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Glucose Intolerance](#)
- [Glucose Metabolism Disorders](#)
- [Hypercholesterolemia](#)
- [Hyperinsulinism](#)
- [Hypoglycemia](#)
- [Insulin Resistance](#)
- [Kidney Failure](#)
- [Metabolic Syndrome X](#)
- [Neoplasms](#)
- [Obesity](#)
- [Osteoporosis](#)
- [Ovarian Failure](#)
- [Overweight](#)
- [Persistent Hyperinsulinemia Hypoglycemia of Infancy](#)
- [Polycystic Ovary Syndrome](#)
- [Puberty](#)
- [Renal Insufficiency](#)
- [Thrombophilia](#)
- [Tobacco Use Disorder](#)
- [Weight Loss](#)
- [Werner syndrome](#)