

## TBL1X (Human) IP-WB Antibody Pair

Catalog # H00006907-PW1 Size 1 Set

## **Applications**



Immunoprecipitation of TBL1X transfected lysate using rabbit polyclonal anti-TBL1X and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with mouse purified polyclonal anti-TBL1X.

Specification	
Product Description	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.
Reactivity	Human
Interspecies Antigen Sequence	Mouse (94); Rat (94)
Quality Control Testing	Immunoprecipitation-Western Blot (IP-WB) Immunoprecipitation of TBL1X transfected lysate using rabbit polyclonal anti-TBL1X and Protein A M agnetic Bead ( <u>U0007</u> ), and immunoblotted with mouse purified polyclonal anti-TBL1X.
Supplied Product	Antibody pair set content:  1. Antibody pair for IP: rabbit polyclonal anti-TBL1X (300 ul)  2. Antibody pair for WB: mouse purified polyclonal anti-TBL1X (50 ug)
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze tha w cycle. Reagents should be returned to -20°C storage immediately after use.

## **Applications**



• Immunoprecipitation-Western Blot

Protocol Download

Gene Info — TBL1X	
Entrez GenelD	<u>6907</u>
Gene Name	TBL1X
Gene Alias	EBI, SMAP55, TBL1
Gene Description	transducin (beta)-like 1X-linked
Omim ID	300196
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene has sequence similarity with members of the WD40 repeat-con taining protein family. The WD40 group is a large family of proteins, which appear to have a regul atory function. It is believed that the WD40 repeats mediate protein-protein interactions and members of the family are involved in signal transduction, RNA processing, gene regulation, vesicular trafficking, cytoskeletal assembly and may play a role in the control of cytotypic differentiation. This encoded protein is found as a subunit in corepressor SMRT (silencing mediator for retinoid and thyroid receptors) complex along with histone deacetylase 3 protein. This gene is located adjacent to the ocular albinism gene and it is thought to be involved in the pathogenesis of the ocular albinism with late-onset sensorineural deafness phenotype. Four transcript variants encoding two different isoforms have been found for this gene. This gene is highly similar to the Y chromosome TBL1Y gene. [provided by RefSeq
Other Designations	F-box-like/WD repeat-containing protein TBL1X OTTHUMP00000022880 transducin beta-like 1X

## Pathway

Wnt signaling pathway