

YARS rabbit monoclonal antibody

Catalog # H00008565-K

Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human YARS peptide using ARM Technology.
Immunogen	A synthetic peptide of human YARS is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human YARS peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — YARS

Entrez GeneID [8565](#)

GeneBank Accession# [YARS](#)

Gene Name YARS

Gene Alias CMTDIC, TYRRS, YRS, YTS

Gene Description tyrosyl-tRNA synthetase

Omim ID [603623 608323](#)

Gene Ontology [Hyperlink](#)

Gene Summary

Aminoacyl-tRNA synthetases catalyze the aminoacylation of tRNA by their cognate amino acid. Because of their central role in linking amino acids with nucleotide triplets contained in tRNAs, aminoacyl-tRNA synthetases are thought to be among the first proteins that appeared in evolution. Tyrosyl-tRNA synthetase belongs to the class I tRNA synthetase family. Cytokine activities have also been observed for the human tyrosyl-tRNA synthetase, after it is split into two parts, an N-terminal fragment that harbors the catalytic site and a C-terminal fragment found only in the mammalian enzyme. The N-terminal fragment is an interleukin-8-like cytokine, whereas the released C-terminal fragment is an EMAP II-like cytokine. [provided by RefSeq]

Other Designations OTTHUMP00000004027|tyrosine tRNA ligase 1, cytoplasmic

Pathway

- [Aminoacyl-tRNA biosynthesis](#)