

PABPC4 rabbit monoclonal antibody

Catalog # H00008761-K

Size 100 ug x up to 3

Specification

| | |
|--------------------------------|--|
| Product Description | Rabbit monoclonal antibody raised against a human PABPC4 peptide using ARM Technology. |
| Immunogen | A synthetic peptide of human PABPC4 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence. |
| Host | Rabbit |
| Library Construction | Non-fusion antibody library from rabbit spleen (ARM Technology). |
| Expression | Overexpression vector and transfection into 293H cell line. |
| Reactivity | Human |
| Purification | Protein A |
| Isotype | IgG |
| Quality Control Testing | Antibody reactive against human PABPC4 peptide by ELISA and mammalian transfected lysate by Western Blot. |
| Storage Buffer | In 1x PBS, pH 7.4 |
| Storage Instruction | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. |
| Deliverable | Up to three rabbit IgG clones of 100 ug each will be delivered to customer. |
| Note | 1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request. |

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — PABPC4

| | |
|---------------------|---|
| Entrez GeneID | 8761 |
| GeneBank Accession# | PABPC4 |
| Gene Name | PABPC4 |
| Gene Alias | APP-1, APP1, FLJ43938, PABP4, iPABP |
| Gene Description | poly(A) binding protein, cytoplasmic 4 (inducible form) |
| Omim ID | 603407 |
| Gene Ontology | Hyperlink |
| Gene Summary | <p>Poly(A)-binding proteins (PABPs) bind to the poly(A) tail present at the 3-prime ends of most eukaryotic mRNAs. PABPC4 or iPABP (inducible PABP) was isolated as an activation-induced T-cell mRNA encoding a protein. Activation of T cells increased PABPC4 mRNA levels in T cells approximately 5-fold. PABPC4 contains 4 RNA-binding domains and proline-rich C terminus. PABPC4 is localized primarily to the cytoplasm. It is suggested that PABPC4 might be necessary for regulation of stability of labile mRNA species in activated T cells. PABPC4 was also identified as an antigen, APP1 (activated-platelet protein-1), expressed on thrombin-activated rabbit platelets. PABPC4 may also be involved in the regulation of protein translation in platelets and megakaryocytes or may participate in the binding or stabilization of polyadenylates in platelet dense granules. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]</p> |
| Other Designations | OTTHUMP00000010668 inducible poly(A)-binding protein poly A binding protein, cytoplasmic 4 |