


Full-Length

CCNE2 (Human) Recombinant Protein (P01)

Catalog # H00009134-P01

Size 10 ug, 25 ug

Applications



Specification

Product Description

Human CCNE2 full-length ORF (AAH20729, 1 a.a. - 374 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence

MSRRSSRLQAKQQPQPSQTESPQEAQIIQAKKRKTTQDVKKRREEVTKKHQYEIRNCWPPVLSG
GISPCIIETPHKEIGTSDFSRFTNYRFKNLFINPSPLPDLSWGCSKEVWLNMLKKESRYVHDKHFE
VLHSDLEPQMRSILLDWLLEVCEVYTLHRETFYLAQDFFDRFMLTQKDINKNMLQLIGITSLFIASKL
EEIYAPKLQEFAYVTDGACSEEDILRMELIILKALKWELCPVTIISWLNFLQVDALKDAPKVLLPQY
SQETFIQIAQLLDLCILAIDSLEFQYRILTAAALCHFTSIEVVKKASGLEWDSISECVDWMVPFVNVV
KSTSPVKLKTFFKIPMEDRHNIQTHTNYLAMLCMISSHV

Host

Wheat Germ (in vitro)

Theoretical MW (kDa)

66.88

Interspecies Antigen Sequence

Mouse (92); Rat (92)

Preparation Method

[in vitro wheat germ expression system](#)

Purification

Glutathione Sepharose 4 Fast Flow

Quality Control Testing

12.5% SDS-PAGE Stained with Coomassie Blue.

Storage Buffer

50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — CCNE2

Entrez GeneID[9134](#)**GeneBank Accession#**[BC020729](#)**Protein Accession#**[AAH20729](#)**Gene Name**

CCNE2

Gene Alias

CYCE2

Gene Description

cyclin E2

Omim ID[603775](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2. This cyclin has been shown to specifically interact with CIP/KIP family of CDK inhibitors, and plays a role in cell cycle G1/S transition. The expression of this gene peaks at the G1-S phase and exhibits a pattern of tissue specificity distinct from that of cyclin E1. A significantly increased expression level of this gene was observed in tumor-derived cells. [provided by RefSeq]

Other Designations

G1/S-specific cyclin E2

Pathway

- [Cell cycle](#)
- [p53 signaling pathway](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Small cell lung cancer](#)

Disease

- [Genetic Predisposition to Disease](#)
- [Ovarian Neoplasms](#)