


ABCC5 (Human) Recombinant Protein (Q01)

Catalog # H00010057-Q01 Size 25 ug, 10 ug

Applications



Specification	
Product Description	Human ABCC5 partial ORF (NP_005679.2, 760 a.a 846 a.a.) recombinant protein with GST-tag a t N-terminal.
Sequence	CITERGTHEELMNLNGDYATIFNNLLLGETPPVEINSKKETSGSQKKSQDKGPKTGSVKKEKAVKP EEGQLVQLEEKGQGSVPWSVY
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	35.31
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications



- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — ABCC5	
Entrez GenelD	<u>10057</u>
GeneBank Accession#	NM_005688
Protein Accession#	NP_005679.2
Gene Name	ABCC5
Gene Alias	ABC33, DKFZp686C1782, EST277145, MOAT-C, MOATC, MRP5, SMRP, pABC11
Gene Description	ATP-binding cassette, sub-family C (CFTR/MRP), member 5
Omim ID	<u>605251</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membrane s. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MRP subfamily which is involved in multi-drug res istance. This protein functions in the cellular export of its substrate, cyclic nucleotides. This export contributes to the degradation of phosphodiesterases and possibly an elimination pathway for cyclic nucleotides. Studies show that this protein provides resistance to thiopurine anticancer drugs, 6-mercatopurine and thioguanine, and the anti-HIV drug 9-(2-phosphonylmethoxyethyl)adenine. This protein may be involved in resistance to thiopurines in acute lymphoblastic leukemia and antiret roviral nucleoside analogs in HIV-infected patients. Alternative splicing of this gene has been detected; however, the complete sequence and translation initiation site is unclear. [provided by RefS eq
Other Designations	ATP-binding cassette, sub-family C, member 5 canalicular multispecific organic anion transporter C

Pathway



ABC transporters

Disease

- Cardiovascular Diseases
- Diabetes Mellitus
- Edema
- Hearing Loss
- Kidney Failure