QPRT rabbit monoclonal antibody

Catalog # H00023475-K

ocification

Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human QPRT peptide using ARM Technology.
Immunogen	A synthetic peptide of human QPRT is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
lsotype	lgG
Quality Control Testing	Antibody reactive against human QPRT peptide by ELISA and mammalian transfected lysate by We stern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

• Western Blot (Transfected lysate)

Protocol Download



• ELISA

Gene Info — QPRT

Entrez GenelD	23475
GeneBank Accession#	<u>QPRT</u>
Gene Name	QPRT
Gene Alias	QPRTase
Gene Description	quinolinate phosphoribosyltransferase
Omim ID	<u>606248</u>
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a key enzyme in catabolism of quinolinate, an intermediate in the tryptophan-n icotinamide adenine dinucleotide pathway. Quinolinate acts as a most potent endogenous exitoto xin to neurons. Elevation of quinolinate levels in the brain has been linked to the pathogenesis of n eurodegenerative disorders such as epilepsy, Alzheimer's disease, and Huntington's disease. [pr ovided by RefSeq
Other Designations	nicotinate-nucleotide pyrophosphorylase (carboxylating)

Pathway

- Biosynthesis of alkaloids derived from ornithine
- <u>Metabolic pathways</u>
- Nicotinate and nicotinamide metabolism