

## ANGPTL4 rabbit monoclonal antibody

Catalog # H00051129-K Size 100 ug x up to 3

| Western Blot.  Storage Buffer In 1x PBS, pH 7.4  Storage Instruction Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.  Deliverable Up to three rabbit IgG clones of 100 ug each will be delivered to customer.  Note 1. Customer may provide cell or tissue lysate for antibody screening.   |                         |  |
|--|-------------------------|--|
| Immunogen       A synthetic peptide of human ANGPTL4 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.         Host       Rabbit         Library Construction       Non-fusion antibody library from rabbit spleen (ARM Technology).         Expression       Overexpression vector and transfection into 293H cell line.         Reactivity       Human         Purification       Protein A         Isotype       IgG         Quality Control Testing       Antibody reactive against human ANGPTL4 peptide by ELISA and mammalian transfected lysate by Western Blot.         Storage Buffer       In 1x PBS, pH 7.4         Storage Instruction       Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.         Deliverable       Up to three rabbit IgG clones of 100 ug each will be delivered to customer.         Note       1. Customer may provide cell or tissue lysate for antibody screening. | Specification           |  |
| Customer or Abnova will decide on the preferred peptide sequence.  Host Rabbit  Library Construction Non-fusion antibody library from rabbit spleen (ARM Technology).  Expression Overexpression vector and transfection into 293H cell line.  Reactivity Human  Purification Protein A  Isotype IgG  Quality Control Testing Antibody reactive against human ANGPTL4 peptide by ELISA and mammalian transfected lysate by Western Blot.  Storage Buffer In 1x PBS, pH 7.4  Storage Instruction Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.  Deliverable Up to three rabbit IgG clones of 100 ug each will be delivered to customer.  Note 1. Customer may provide cell or tissue lysate for antibody screening.  | Product Description     | Rabbit monoclonal antibody raised against a human ANGPTL4 peptide using ARM Technology.                    |
| Library Construction  Non-fusion antibody library from rabbit spleen (ARM Technology).  Expression  Overexpression vector and transfection into 293H cell line.  Reactivity  Human  Purification  Protein A  Isotype  IgG  Quality Control Testing  Antibody reactive against human ANGPTL4 peptide by ELISA and mammalian transfected lysate by Western Blot.  Storage Buffer  In 1x PBS, pH 7.4  Storage Instruction  Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.  Deliverable  Up to three rabbit IgG clones of 100 ug each will be delivered to customer.  Note  1. Customer may provide cell or tissue lysate for antibody screening.  | Immunogen               |  |
| Expression Overexpression vector and transfection into 293H cell line.  Reactivity Human  Purification Protein A  Isotype IgG  Quality Control Testing Antibody reactive against human ANGPTL4 peptide by ELISA and mammalian transfected lysate by Western Blot.  Storage Buffer In 1x PBS, pH 7.4  Storage Instruction Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.  Deliverable Up to three rabbit IgG clones of 100 ug each will be delivered to customer.  Note 1. Customer may provide cell or tissue lysate for antibody screening.   | Host                    | Rabbit   |
| Reactivity       Human         Purification       Protein A         Isotype       IgG         Quality Control Testing       Antibody reactive against human ANGPTL4 peptide by ELISA and mammalian transfected lysate by Western Blot.         Storage Buffer       In 1x PBS, pH 7.4         Storage Instruction       Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.         Deliverable       Up to three rabbit IgG clones of 100 ug each will be delivered to customer.         Note       1. Customer may provide cell or tissue lysate for antibody screening.  | Library Construction    | Non-fusion antibody library from rabbit spleen ( <u>ARM Technology</u> ).                                  |
| Purification Protein A  Isotype IgG  Quality Control Testing Antibody reactive against human ANGPTL4 peptide by ELISA and mammalian transfected lysate by Western Blot.  Storage Buffer In 1x PBS, pH 7.4  Storage Instruction Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.  Deliverable Up to three rabbit IgG clones of 100 ug each will be delivered to customer.  Note 1. Customer may provide cell or tissue lysate for antibody screening.   | Expression              | Overexpression vector and transfection into 293H cell line.  |
| Isotype   IgG  | Reactivity              | Human  |
| Quality Control Testing       Antibody reactive against human ANGPTL4 peptide by ELISA and mammalian transfected lysate by Western Blot.         Storage Buffer       In 1x PBS, pH 7.4         Storage Instruction       Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.         Deliverable       Up to three rabbit lgG clones of 100 ug each will be delivered to customer.         Note       1. Customer may provide cell or tissue lysate for antibody screening.  | Purification            | Protein A  |
| Western Blot.  Storage Buffer In 1x PBS, pH 7.4  Storage Instruction Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.  Deliverable Up to three rabbit IgG clones of 100 ug each will be delivered to customer.  Note 1. Customer may provide cell or tissue lysate for antibody screening.   | Isotype                 | lgG  |
| Storage Instruction       Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.         Deliverable       Up to three rabbit lgG clones of 100 ug each will be delivered to customer.         Note       1. Customer may provide cell or tissue lysate for antibody screening.  | Quality Control Testing | Antibody reactive against human ANGPTL4 peptide by ELISA and mammalian transfected lysate by Western Blot. |
| Deliverable  Up to three rabbit lgG clones of 100 ug each will be delivered to customer.  Note  1. Customer may provide cell or tissue lysate for antibody screening.  | Storage Buffer          | In 1x PBS, pH 7.4  |
| Note 1. Customer may provide cell or tissue lysate for antibody screening.   | Storage Instruction     | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.                                   |
| o action in the promise of a least of an arrange   | Deliverable             | Up to three rabbit IgG clones of 100 ug each will be delivered to customer.                                |
| cluding F(ab) <sub>2</sub> , lgG, scFv and different Fc and non-Fc conjugates per customer request.  | Note                    | 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in           |

## **Applications**

Western Blot (Transfected lysate)

Protocol Download



ELISA

| Gene Info — ANGPTL4 |   |
|---------------------|---|
| Entrez GenelD       | <u>51129</u>  |
| GeneBank Accession# | ANGPTL4   |
| Gene Name           | ANGPTL4   |
| Gene Alias          | ANGPTL2, ARP4, FIAF, HFARP, NL2, PGAR, pp1158   |
| Gene Description    | angiopoietin-like 4   |
| Omim ID             | 605910  |
| Gene Ontology       | <u>Hyperlink</u>  |
| Gene Summary        | This gene is a member of the angiopoietin/angiopoietin-like gene family and encodes a glycosyla ted, secreted protein with a fibrinogen C-terminal domain. This gene is induced under hypoxic conditions in endothelial cells and is the target of peroxisome proliferation activators. The encoded protein is a serum hormone directly involved in regulating glucose homeostasis, lipid metabolism, and insulin sensitivity and also acts as an apoptosis survival factor for vascular endothelial cells. The encoded protein may play a role in several cancers and it also has been shown to prevent the metastatic process by inhibiting vascular activity as well as tumor cell motility and invasiveness. Decreased expression of this protein has been associated with type 2 diabetes. Alternatively splice d transcript variants encoding different isoforms have been described. This gene was previously referred to as ANGPTL2 but has been renamed ANGPTL4. [provided by RefSeq |
| Other Designations  | PPARG angiopoietin related protein angiopoietin-like 4 protein angiopoietin-related protein 4 fast ing-induced adipose factor hepatic angiopoietin-related protein hepatic fibrinogen/angiopoietin-related protein peroxisome proliferator-activated receptor (P  |

## Pathway

PPAR signaling pathway

## Disease

- Amyotrophic lateral sclerosis
- Anoxia
- Atherosclerosis



- Cardiovascular Diseases
- Coronary Artery Disease
- Coronary Disease
- Diabetes Mellitus
- Dyslipidemias
- Edema
- Genetic Predisposition to Disease
- Kidney Failure
- Syndrome
- Weight Gain