


MaxPab®

TNFRSF19 purified MaxPab mouse polyclonal antibody (B01P)

Catalog # H00055504-B01P

Size 50 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of TNFRSF19 expression in transfected 293T cell line ([H00055504-T01](#)) by TNFRSF19 MaxPab polyclonal antibody.

Lane 1: TNFRSF19 transfected lysate(45.30 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description	Mouse polyclonal antibody raised against a full-length human TNFRSF19 protein.
Immunogen	TNFRSF19 (NP_683760.1, 1 a.a. ~ 417 a.a) full-length human protein.
Sequence	MALKVLLQEKTFFLLVLLGYLSCKVTCESGDCRQQEFRDRSGNCVPCNQCGPMELSKECG FGYGEDAQCVTCLHRFKEDWGFQKCKPCLDCAVVNRFQKANCSATSDAICGDCLPGFYRKT LVGFQDMECVPCGDPPIYEPHCASKVNLVKIASTASSPRDTALAAVICSALATVLLALLILCVYC KRQFMEKKPSWSLRSQDIQYNGSELSCFDRPQLHEYAHRACCQCRSDSVQTCGPVRLLP SMC CEEACSPNPATLGCVHSAASLQARNAGPAGEMVPTFFGSLTQSICGEFSDAWPLMQNPMGG DNISFCDSPELTGEDIHSLNPELESSTSLDSNSSQDLVGGAVPVQSHSENFTAATDL SRYNNTLV ESASTQDALTMRSQLDQESGAVIHPATQTSLQEA
Host	Mouse
Reactivity	Human
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of TNFRSF19 expression in transfected 293T cell line ([H00055504-T01](#)) by TNFRSF19 MaxPab polyclonal antibody.

Lane 1: TNFRSF19 transfected lysate(45.30 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — TNFRSF19

Entrez GeneID [55504](#)

GeneBank Accession# [NM_148957](#)

Protein Accession# [NP_683760.1](#)

Gene Name TNFRSF19

Gene Alias TAJ, TAJ-alpha, TRADE, TROY

Gene Description tumor necrosis factor receptor superfamily, member 19

Omim ID [606122](#)

Gene Ontology [Hyperlink](#)

Gene Summary The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is highly expressed during embryonic development. It has been shown to interact with TRAF family members, and to activate JNK signaling pathway when overexpressed in cells. This receptor is capable of inducing apoptosis by a caspase-independent mechanism, and it is thought to play an essential role in embryonic development. Alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]

Other Designations OTTHUMP00000018113|OTTHUMP00000018114|toxicity and JNK inducer

Pathway

- [Cytokine-cytokine receptor interaction](#)

Disease

- [Asthma](#)
- [Genetic Predisposition to Disease](#)
- [Nasopharyngeal Neoplasms](#)