

PCDHA8 rabbit monoclonal antibody

Catalog # H00056140-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human PCDHA8 peptide using ARM Technology.
Immunogen	A synthetic peptide of human PCDHA8 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human PCDHA8 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — PCDHA8

Entrez GeneID [56140](#)

GeneBank Accession# [PCDHA8](#)

Gene Name PCDHA8

Gene Alias PCDH-ALPHA8

Gene Description protocadherin alpha 8

Omim ID [606314](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene is a member of the protocadherin alpha gene cluster, one of three related gene clusters tandemly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related to the mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding sequences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-terminal exons, or constant exons, which are shared by all genes in the cluster. The large, uninterrupted N-terminal exons each encode six cadherin ectodomains while the C-terminal exons encode the cytoplasmic domain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been observed and additional variants have been suggested but their full-length nature has yet to be determined. [provided by RefSeq]

Other Designations KIAA0345-like 6