


NOL6 293T Cell Transient Overexpression Lysate(Denatured)


Catalog # H00065083-T01 Size 100 uL

Applications



SDS-PAGE Gel

NOL6 transfected lysate.



Western Blot

Lane 1: NOL6 transfected lysate (22.11 KDa)

Lane 2: Non-transfected lysate.

Specification	
Transfected Cell Line	293T
Plasmid	pCMV-NOL6 full-length
Host	Human
Theoretical MW (kDa)	22.11
Interspecies Antigen Sequence	Mouse (84); Rat (86)



Product Information

Quality Control Testing	Transient overexpression cell lysate was tested with Anti-NOL6 antibody (H00065083-B01) by West ern Blots. SDS-PAGE Gel NOL6 transfected lysate. Western Blot Lane 1: NOL6 transfected lysate (22.11 KDa) Lane 2: Non-transfected lysate.
Storage Buffer	1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue)
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

Western Blot

Gene Info — NOL6	
Entrez GenelD	<u>65083</u>
GeneBank Accession#	BC008298
Protein Accession#	AAH08298
Gene Name	NOL6
Gene Alias	FLJ21959, MGC14896, MGC14921, MGC20838, NRAP, UTP22, bA311H10.1
Gene Description	nucleolar protein family 6 (RNA-associated)
Omim ID	611532
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The nucleolus is a dense subnuclear membraneless organelle that assembles around clusters of r RNA genes and functions in ribosome biogenesis. This gene encodes a nucleolar RNA-associat ed protein that is highly conserved between species. RNase treatment of permeabilized cells indicates that the nucleolar localization is RNA dependent. Further studies suggest that the protein is associated with ribosome biogenesis through an interaction with pre-rRNA primary transcripts. Alt ernative splicing has been observed at this locus and two splice variants encoding distinct isoforms have been identified. [provided by RefSeq
Other Designations	OTTHUMP0000000449 OTTHUMP0000000450 OTTHUMP0000000451 OTTHUMP000000000000000000000000000000000000