TNFRSF8 monoclonal antibody, clone CD30/412

Catalog # MAB11230 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of Hodgkin's lymphoma using TNFRSF8 monoclonal antibody, clone CD30/412 (Cat # MAB11230).

Specification

Product Description	Mouse monoclonal antibody raised against partial recombinant TNFRSF8.
Immunogen	Recombinant protein corresponding to human TNFRSF8.
Host	Mouse
Reactivity	Human
Form	Liquid
Purification	Protein G affinity chromatography
Isotype	IgG1, kappa
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:200-1:400) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% BSA, 0.05% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of Hodgkin's lymphoma using TNFRSF8 monoclonal antibody, clone CD30/412 (Cat # MAB11230).

Gene Info — TNFRSF8

Entrez GeneID	943
Protein Accession#	NP_001234
Gene Name	TNFRSF8
Gene Alias	CD30, D1S166E, KI-1
Gene Description	tumor necrosis factor receptor superfamily, member 8
Omim ID	153243
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This rece ptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potenti al of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternativel y spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq
Other Designations	CD30 antigen|CD30L receptor|Ki-1 antigen|OTTHUMP00000001783|cytokine receptor CD30|ly mphocyte activation antigen CD30

Pathway

- Cytokine-cytokine receptor interaction

Disease

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Asthma
- Diabetes Mellitus
- Genetic Predisposition to Disease
- Hematologic Diseases
- HIV Infections
- Kidney Failure
- Multiple Myeloma
- Occupational Diseases
- Tobacco Use Disorder

Copyright © 2023 Abnova Corporation. All Rights Reserved.