
BCL2 monoclonal antibody, clone 8C8


Catalog # MAB11265 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human tonsil with BCL2 monoclonal antibody, clone 8C8 (Cat # MAB11265) at 1:200 dilution.



Western Blot (Cell lysate)

Western blot analysis of cell lysates by BCL2 monoclonal antibody, clone 8C8 (Cat # MAB11265) at 2 ug/ml. Lane1 : HeLa. Lane2 : A549. Lane3 : MCF7.

Specification

Product Description	Mouse monoclonal antibody raised against synthetic peptide of BCL2.
Immunogen	A synthetic peptide corresponding to amino acids 41-54 of human BCL2.
Sequence	GAAPAPGIFSSQPG
Host	Mouse
Reactivity	Human, Monkey
Form	Liquid

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Purification	Protein G purification
Isotype	IgG1 Kappa
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:200-1:400) Western Blot (1-3 ug/ml) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% BSA, 0.05% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical staining of human tonsil with BCL2 monoclonal antibody, clone 8C8 (Cat # MAB11265) at 1:200 dilution.

- Immunofluorescence
- Immunocytochemistry
- Flow Cytometry
- Western Blot (Cell lysate)

  Western blot analysis of cell lysates by BCL2 monoclonal antibody, clone 8C8 (Cat # MAB11265) at 2 ug/ml. Lane1 : HeLa. Lane2 : A549. Lane3 : MCF7.

Gene Info — BCL2

Entrez GeneID	596
Protein Accession#	NP_000624
Gene Name	BCL2
Gene Alias	Bcl-2
Gene Description	B-cell CLL/lymphoma 2
Omim ID	151430

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Gene Ontology	Hyperlink
Gene Summary	This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends. [provided by RefSeq
Other Designations	B-cell lymphoma protein 2|OTTHUMP00000163680

Pathway

- Amyotrophic lateral sclerosis (ALS)
- Apoptosis
- Colorectal cancer
- Focal adhesion
- Neurotrophin signaling pathway
- Pathways in cancer
- Prostate cancer
- Small cell lung cancer

Disease

- Adenocarcinoma
- Alzheimer Disease
- Amnesia
- Attention Deficit Disorder with Hyperactivity
- Autistic Disorder
- Azoospermia
- Bile Duct Neoplasms
- Brain Injuries
- Breast cancer

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Breast Neoplasms
- Carcinoma
- Cardiovascular Diseases
- Cholangiocarcinoma
- Chromosome Aberrations
- Clubfoot
- Cognition
- Cognition Disorders
- Diabetes Mellitus
- Disease Progression
- Edema
- Esophageal Neoplasms
- Genetic Predisposition to Disease
- Head and Neck Neoplasms
- Hematologic Diseases
- Hodgkin Disease
- Kidney Failure
- Kidney Neoplasms
- Laryngeal Neoplasms
- Leukemia
- Leukemia-Lymphoma
- Lung Neoplasms
- Lupus Erythematosus
- Lymphatic Metastasis
- Lymphoma

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Lymphoproliferative Disorders
- Multiple Sclerosis
- NARP
- Neoplasm Invasiveness
- Neoplasm Recurrence
- Neovascularization
- Neuropsychological Tests
- Occupational Diseases
- Oropharyngeal Neoplasms
- Ovarian Neoplasms
- Postoperative Complications
- Prostate cancer
- Prostatic Neoplasms
- Prosthesis Failure
- Pulmonary Disease
- Thyroid Neoplasms
- Translocation
- Waldenstrom Macroglobulinemia
- Werner syndrome

Copyright © 2023 Abnova Corporation. All Rights Reserved.