
CD8A monoclonal antibody, clone RPA-T8 (FITC)

Catalog # MAB12213 Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of human peripheral blood lymphocytes using CD8A monoclonal antibody, clone RPA-T8 (FITC) (Cat # MAB12213) compared to a relevant isotype control in red.

| Specification | |
|---------------------|--|
| Product Description | Mouse monoclonal antibody raised against CD8A. |
| Immunogen | CD8A |
| Host | Mouse |
| Reactivity | Human |
| Form | Liquid |
| Conjugation | FITC |
| Purification | Affinity Chromatography |
| lsotype | lgG1, kappa |
| Recommend Usage | Flow Cytometry The optimal working dilution should be determined by the end user. |
| Storage Buffer | In PBS, pH7.2 (0.09% Sodium azide, may contain carrier protein/stabilizer). |
| Storage Instruction | Store in the dark at 4°C. Avoid prolonged exposure to light. Do not freeze. |

Copyright © 2023 Abnova Corporation. All Rights Reserved.

😵 Abnova

Product Information

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Flow Cytometry

Flow cytometric analysis of human peripheral blood lymphocytes using CD8A monoclonal antibody, clone RPA-T8 (FITC) (Cat # MAB12213) compared to a relevant isotype control in red.

| Gene Info — CD8A | |
|--------------------|---|
| Entrez GenelD | <u>925</u> |
| Gene Name | CD8A |
| Gene Alias | CD8, Leu2, MAL, p32 |
| Gene Description | CD8a molecule |
| Omim ID | <u>186910 608957</u> |
| Gene Ontology | Hyperlink |
| Gene Summary | The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediat es efficient cell-cell interactions within the immune system. The CD8 antigen acts as a corepresso r with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen prese nting cell (APC) in the context of class I MHC molecules. The coreceptor functions as either a hom odimer composed of two alpha chains, or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light cha ins. This gene encodes the CD8 alpha chain isoforms. Multiple transcript variants encoding differ ent isoforms have been found for this gene. [provided by RefSeq |
| Other Designations | CD8 antigen alpha polypeptide CD8 antigen, alpha polypeptide (p32) Leu2 T-lymphocyte antigen OKT8 T-cell antigen T cell co-receptor T-cell antigen Leu2 T-cell surface glycoprotein CD8 alpha c hain T-lymphocyte differentiation antigen T8/Leu-2 T8 T-cell ant |

Pathway

- Antigen processing and presentation
- <u>Cell adhesion molecules (CAMs)</u>
- Hematopoietic cell lineage

😵 Abnova

- Primary immunodeficiency
- <u>T cell receptor signaling pathway</u>