




# Histone H3 (acetyl K36) monoclonal antibody, clone RM154


Catalog # MAB12836 Size 100 ug

## **Applications**



### Western Blot (Cell lysate)

Western blot analysis of Lane 1: HeLa cells, Lane 2: HeLa cells treated with sodium butyrate using Histone H3 (acetyl K36) monoclonal antibody, clone RM154 (Cat # MAB12836) under 1 ug/mL working concentration.



#### **Immunocytochemistry**

Immunocytochemistry staining of HeLa cells treated with sodium butyrate using Histone H3 (acetyl K36) monoclonal antibody, clone RM154 (Cat # MAB12836) (Red) under 2 ug/mL working concentration. Actin filaments were stained with fluorescein phalloidin (Green).

Specification	
Product Description	Rabbit recombinant monoclonal antibody raised against of human histone H3 (acetyl K36).
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic acetyl peptide corresponding to residues surrounding K36 of human Histone H3.
Sequence	N/A



#### **Product Information**

Specificity	This antibody reacts to histone H3 acetylated at Lysine 36, and is not affected by the phosphorylation of neighboring Ser31. No cross reactivity with other acetylated Lysines in histone H3.
Form	Liquid
Purification	Protein A affinity purification
Isotype	lgG
Recommend Usage	ELISA (0.2-1 ug/mL) Immunocytochemistry (0.5-2 ug/mL) Western Blot (0.5-2 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## **Applications**

Western Blot (Cell lysate)

Western blot analysis of Lane 1: HeLa cells, Lane 2: HeLa cells treated with sodium butyrate using Histone H3 (acetyl K36) monoclonal antibody, clone RM154 (Cat # MAB12836) under 1 ug/mL working concentration.

Immunocytochemistry

Immunocytochemistry staining of HeLa cells treated with sodium butyrate using Histone H3 (acetyl K36) monoclonal antibody, clone RM154 (Cat # MAB12836) (Red) under 2 ug/mL working concentration. Actin filaments were stained with fluorescein phalloidin (Green).

Enzyme-linked Immunoabsorbent Assay

## Gene Info — HIST1H3A

Entrez GeneID	<u>8350</u>
Gene Name	HIST1H3A
Gene Alias	H3/A, H3FA
Gene Description	histone cluster 1, H3a



## **Product Information**

Omim ID	<u>602810</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chro mosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped ar ound a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H 1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA t ails; instead, they contain a palindromic termination element. This gene is found in the large histon e gene cluster on chromosome 6p22-p21.3. [provided by RefSeq
Other Designations	H3 histone family, member A histone 1, H3a

# Pathway

• Systemic lupus erythematosus